



# KCNH6 Polyclonal Antibody

<b>Catalog No</b>	BYab-05957
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	KCNH6 ERG2
<b>Protein Name</b>	Potassium voltage-gated channel subfamily H member 6 (Ether-a-go-go-related gene potassium channel 2) (ERG-2) (Eag-related protein 2) (Ether-a-go-go-related protein 2) (hERG-2) (hERG2) (Voltage-gated
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 230-310
<b>Specificity</b>	KCNH6 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	109kD
<b>Cell Pathway</b>	Membrane; Multi-pass membrane protein.
<b>Tissue Specificity</b>	Expressed in prolactin-secreting adenomas.
<b>Function</b>	alternative products:Experimental confirmation may be lacking for some isoforms, domain:The segment S4 is probably the voltage-sensor and is characterized by a series of positively charged amino acids at every third position., function:Pore-forming (alpha) subunit of voltage-gated potassium channel. Elicits a slowly activating, rectifying current (By similarity). Channel properties may be modulated by cAMP and subunit assembly., similarity:Belongs to the potassium channel family. H (Eag) subfamily., similarity:Contains 1 cyclic nucleotide-binding domain., similarity:Contains 1 PAC (PAS-associated C-terminal) domain., similarity:Contains 1 PAS (PER-ARNT-SIM) domain., subunit:The potassium channel is probably composed of a homo- or heterotetrameric complex of pore-forming alpha subunits that can associate with

Nanjing BYabscience technology Co.,Ltd



modulating beta subunits. Heteromultimers with KCNH2/ERG1 and KCNH7/ERG3.,tissue specif

**Background**

Voltage-gated potassium (Kv) channels represent the most complex class of voltage-gated ion channels from both functional and structural standpoints. Their diverse functions include regulating neurotransmitter release, heart rate, insulin secretion, neuronal excitability, epithelial electrolyte transport, smooth muscle contraction, and cell volume. This gene encodes a member of the potassium channel, voltage-gated, subfamily H. This member is a pore-forming (alpha) subunit. Alternative splicing results in multiple transcript variants that encode different isoforms. [provided by RefSeq, Jul 2013],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**