



ATP4A Polyclonal Antibody

Catalog No	BYab-05964
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	ATP4A
Protein Name	Potassium-transporting ATPase alpha chain 1 (EC 3.6.3.10) (Gastric H(+)/K(+) ATPase subunit alpha) (Proton pump)
Immunogen	Synthesized peptide derived from human protein . at AA range: 800-880
Specificity	ATP4A Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	113kD
Cell Pathway	Apical cell membrane ; Multi-pass membrane protein . Localized in the apical canalicular membrane of parietal cells (PubMed:24188822). Localized in the apical canalicular membrane of parietal cells (PubMed:24188822). .
Tissue Specificity	Expressed in gastric parietal cells (at protein level).
Function	catalytic activity:ATP + H(2)O + H(+)(In) + K(+)(Out) = ADP + phosphate + H(+)(Out) + K(+)(In).,function:Catalyzes the hydrolysis of ATP coupled with the exchange of H(+) and K(+) ions across the plasma membrane. Responsible for acid production in the stomach.,similarity:Belongs to the cation transport ATPase (P-type) family.,similarity:Belongs to the cation transport ATPase (P-type) family. Type IIC subfamily.,subunit:Composed of two subunits: alpha (catalytic) and beta.,tissue specificity:Found in gastric mucosa.,
Background	The protein encoded by this gene belongs to a family of P-type cation-transporting ATPases. The gastric H+, K+-ATPase is a heterodimer consisting of a high molecular weight catalytic alpha subunit and a smaller but

Nanjing BYabscience technology Co.,Ltd



heavily glycosylated beta subunit. This enzyme is a proton pump that catalyzes the hydrolysis of ATP coupled with the exchange of H(+) and K(+) ions across the plasma membrane. It is also responsible for gastric acid secretion. This gene encodes a catalytic alpha subunit of the gastric H⁺, K⁺-ATPase. [provided by RefSeq, Jul 2008],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images