



PP1RA Polyclonal Antibody

Catalog No BYab-06145 Isotype IgG Reactivity Human; Mouse; Rat Applications WB; ELISA Gene Name PPP1R10 CAT53 FB19 PNUTS Protein Name Serine/threonine-protein phosphatase 1 regulatory subunit 10 (MHC class I region proline-rich protein CAT53) (PP1-binding protein of 114 kDa) (Phosphatase 1 inclear targeting subunit) (Protein FB19) (Immunogen Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidne		
Reactivity Human;Mouse;Rat Applications WB;ELISA Gene Name PPP1R10 CAT53 FB19 PNUTS Protein Name Serine/threonine-protein phosphatase 1 regulatory subunit 10 (MHC class I region proline-rich protein CAT53) (PP1-binding protein of 114 kDa) (Phosphatase 1 nuclear targeting subunit) (Protein FB19) (Immunogen) Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function: Inhibitor of PPP1CA and PPP1CC phosphatase activities	Catalog No	BYab-06145
Applications WB;ELISA Gene Name PPP1R10 CAT53 FB19 PNUTS Protein Name Serine/threonine-protein phosphatase 1 regulatory subunit 10 (MHC class I region proline-rich protein CAT53) (PP1-binding protein of 114 kDa) (Phosphatase 1 nuclear targeting subunit) (Protein FB19) (Immunogen Synthesized peptide derived from part region of human protein Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (sSDNA), poly(A) and poly(G) homopolymers. PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity. Contains 1 TISIS N-terminal domain , subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli. Associates with region necessary for interaction with PPP1CA, similarity. Contains 1 TISIS N-terminal domain , subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli. Associates	Isotype	IgG
Gene Name PPP1R10 CAT53 FB19 PNUTS Protein Name Serine/threonine-protein phosphatase 1 regulatory subunit 10 (MHC class I region proline-rich protein CAT53) (PP1-binding protein of 114 kDa) (Phosphatase 1 nuclear targeting subunit) (Protein FB19) (Immunogen Synthesized peptide derived from part region of human protein Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA, with	Reactivity	Human;Mouse;Rat
Protein Name Serine/threonine-protein phosphatase 1 regulatory subunit 10 (MHC class I region proline-rich protein CAT53) (PP1-binding protein of 114 kDa) (Phosphatase 1 nuclear targeting subunit) (Protein FB19) (Synthesized peptide derived from part region of human protein Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (sSDNA), poly(G) amd poly(G) homopolymers. PTM: Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction within nucleoii, subunit.Interaction within nucleoic plasmic bodies and within nucleoic, subunit.Interaction within nucleoic plasmic bodies and within nucleoic plasmic bodies and within nucleoic plasmic bodies and within nucleoic, subunit.Interaction within nucleoic plasmic bodies and wi	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA and PPP1CC. Issue interaction with PPP1CA and PPP1CC. Sand PPP1CA and PPP1CC. And PPP1CA and PPP1CC. Sand PPP1CA and PPP1CC. Sand PPP	Gene Name	PPP1R10 CAT53 FB19 PNUTS
Specificity PP1RA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers., PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity. Contains 1 C3H1-type zinc finger, similarity. Contains 1 TSIIS N-terminal domain., subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli. subunitic Interacts with PPP1CA and PPP1CC, tissue	Protein Name	Serine/threonine-protein phosphatase 1 regulatory subunit 10 (MHC class I region proline-rich protein CAT53) (PP1-binding protein of 114 kDa) (Phosphatase 1 nuclear targeting subunit) (Protein FB19) (
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers. PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity:Contains 1 C3H1-type zinc finger. similarity:Contains 1 TFIIS N-terminal domain in subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli, subupunit:Interacts with PPP1CA and PPP1CC, dissue	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers. PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA., similarity:Contains 1 C3H1-type zinc finger., similarity:Contains 1 TFIIS N-terminal domain., subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli, subunit:Interacts with PPP1CA and PPP1CC, tissue	Specificity	PP1RA Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleolic, subunit:Interacts with PPP1CA and PPP1CC, tissue	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function: Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers. PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA., similarity:Contains 1 C3H1-type zinc finger., similarity:Contains 1 TFIIS N-terminal domain., subbcellular location:Found in discrete nucleoplasmic bodies and within nucleoli., subunit:Interacts with PPP1CA and PPP1CC., tissue	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC, tissue	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC, tissue	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 103kD Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA, similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC, tissue	Concentration	1 mg/ml
Synonyms Observed Band 103kD Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue	Purity	≥90%
Observed Band Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Found in discrete nucleoplasmic bodies and within nucleoli. Associates with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue	Synonyms	
with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By similarity). Tissue Specificity Expressed in heart, brain, placenta, lung, liver, skeletal muscle, kidney and pancreas. Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue	Observed Band	103kD
Function function:Inhibitor of PPP1CA and PPP1CC phosphatase activities. Has inhibitory activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue	Cell Pathway	with chromatin during interphase, excluded from condensed chromosomes during early mitosis and is reloaded onto chromosomes at the late telophase (By
activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue	Tissue Specificity	
	Function	activity on PPP1CA only when phosphorylated. Binds to mRNA, single-stranded DNA (ssDNA), poly(A) and poly(G) homopolymers.,PTM:Phosphorylated on Ser-398 and Thr-400 by PKA within the region necessary for interaction with PPP1CA.,similarity:Contains 1 C3H1-type zinc finger.,similarity:Contains 1 TFIIS N-terminal domain.,subcellular location:Found in discrete nucleoplasmic bodies and within nucleoli.,subunit:Interacts with PPP1CA and PPP1CC.,tissue

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	and pancreas.,
Background	This gene encodes a protein phosphatase 1 binding protein. The encoded protein plays a role in many cellular processes including cell cycle progression, DNA repair and apoptosis by regulating the activity of protein phosphatase 1. This gene lies within the major histocompatibility complex class I region on chromosome 6, and alternatively spliced transcript variants have been observed for this gene. [provided by RefSeq, Jul 2012],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd