



STAC Polyclonal Antibody

Catalog No	BYab-06168
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	STAC
Protein Name	SH3 and cysteine-rich domain-containing protein (Src homology 3 and cysteine-rich domain-containing protein)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	STAC Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	44kD
Cell Pathway	Cytoplasm, cytosol . Cell membrane ; Peripheral membrane protein ; Cytoplasmic side . Cell membrane, sarcolemma ; Peripheral membrane protein ; Cytoplasmic side .
Tissue Specificity	Brain,Kidney,
Function	function:Probably involved in a neuron-specific signal transduction.,similarity:Contains 1 phorbol-ester/DAG-type zinc finger.,similarity:Contains 1 SH3 domain.,
Background	function:Probably involved in a neuron-specific signal transduction.,similarity:Contains 1 phorbol-ester/DAG-type zinc finger.,similarity:Contains 1 SH3 domain.,
matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

