



SH3K1 Polyclonal Antibody

Catalog No	BYab-06177
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	SH3KBP1 CIN85
Protein Name	SH3 domain-containing kinase-binding protein 1 (CD2-binding protein 3) (CD2BP3) (Cbl-interacting protein of 85 kDa) (Human Src family kinase-binding protein 1) (HSB-1)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	SH3K1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	73kD
Cell Pathway	Cytoplasm . Cytoplasm, cytoskeleton. Cytoplasmic vesicle membrane; Peripheral membrane protein. Cell junction, synapse, synaptosome. Cell junction, focal adhesion . Localized in endocytic vesicles containing clustered receptors. Colocalizes with ASAP1 in vesicular structures. Colocalized with actin microfilaments and focal adhesions (By similarity). Colocalized with MAGI2 in synaptosomes. Translocation to EGFR containing vesicles upon EGF stimulation is inhibited in the presence of SH3KBP1 (By similarity). Colocalizes with ZFP36 in the cytoplasm (PubMed:20221403). .
Tissue Specificity	Ubiquitously expressed. Also expressed in some cancer cell lines.
Function	domain:The SH3 domains mediate interaction with SHKBP1.,function:Adapter protein involved in regulating diverse signal transduction pathways. Involved in the regulation of endocytosis and lysosomal degradation of ligand-induced receptor tyrosine kinases, including EGFR and MET/hepatocyte growth factor

Nanjing BYabscience technology Co.,Ltd



receptor, through a association with CBL and endophilins. The association with CBL, and thus the receptor internalization, may inhibited by an interaction with PDCD6IP and/or SPRY2. Involved in regulation of ligand-dependent endocytosis of the IgE receptor. Attenuates phosphatidylinositol 3-kinase activity by interaction with its regulatory subunit (By similarity). May be involved in regulation of cell adhesion; promotes the interaction between TTK2B and PDCD6IP. May be involved in the regulation of cellular stress response via the MAPK pathways through its interaction with MAP3K4. Is invol

Background

SH3 domain containing kinase binding protein 1(SH3KBP1) Homo sapiens
This gene encodes an adapter protein that contains three N-terminal Src homology domains, a proline rich region and a C-terminal coiled-coil domain. The encoded protein facilitates protein-protein interactions and has been implicated in numerous cellular processes including apoptosis, cytoskeletal rearrangement, cell adhesion and in the regulation of clathrin-dependent endocytosis. Alternate splicing results in multiple transcript variants.[provided by RefSeq, May 2010],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images