



KCNN1 Polyclonal Antibody

| Catalog No | BYab-06186 |
|--------------------|---|
| Isotype | lgG |
| Reactivity | Human;Mouse;Rat |
| Applications | WB;ELISA |
| Gene Name | KCNN1 SK |
| Protein Name | Small conductance calcium-activated potassium channel protein 1 (SK1) (SKCa 1) (SKCa1) (KCa2.1) |
| Immunogen | Synthesized peptide derived from human protein . at AA range: 360-440 |
| Specificity | KCNN1 Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 59kD |
| Cell Pathway | Membrane; Multi-pass membrane protein. |
| Tissue Specificity | Fetal brain,Hippocampus, |
| Function | function:Forms a voltage-independent potassium channel activated by intracellular calcium. Activation is followed by membrane hyperpolarization. Thought to regulate neuronal excitability by contributing to the slow component of synaptic afterhyperpolarization. The channel is blocked by apamin.,similarity:Belongs to the potassium channel KCNN family.,subunit:Heterooligomer. The complex is composed of 4 channel subunits each of which binds to a calmodulin subunit which regulates the channel activity through calcium-binding., |
| Background | potassium calcium-activated channel subfamily N member 1(KCNN1) Homo sapiens Action potentials in vertebrate neurons are followed by an afterhyperpolarization (AHP) that may persist for several seconds and may have |
| | Nanjing BYabscience technology Co.,Ltd |



国内优质抗体供应商

精准的 WB 检测服务 24H 在线服务, 欢迎咨询



| | profound consequences for the firing pattern of the neuron. Each component of the AHP is kinetically distinct and is mediated by different calcium-activated potassium channels. The protein encoded by this gene is activated before membrane hyperpolarization and is thought to regulate neuronal excitability by contributing to the slow component of synaptic AHP. The encoded protein is an integral membrane protein that forms a voltage-independent calcium-activated channel with three other calmodulin-binding subunits. This gene is a member of the KCNN family of potassium channel genes. [provided by RefSeq, Jul 2008], |
|---------------------------|---|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images

Nanjing BYabscience technology Co.,Ltd