



TLX3 Polyclonal Antibody

in children and young adults. [provided by RefSeq, Nov 2015],		
Reactivity Human; Mouse Applications WB; ELISA Gene Name TLX3 HOX11L2 Protein Name T-cell leukemia homeobox protein 3 (Homeobox protein Hox-11L2) Immunogen Synthesized peptide derived from part region of human protein Specificity TLX3 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus. Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5:14)(q35;q32)] involving this gene is associated with T-cell acute lymphostatic leukemia (T-ALL) in children and young adults. [provided by RefSeq, Nov 2015],	Catalog No	BYab-06271
Applications Gene Name TLX3 HOX11L2 Protein Name T-cell leukemia homeobox protein 3 (Homeobox protein Hox-11L2) Immunogen Synthesized peptide derived from part region of human protein Specificity TLX3 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function Similarity:Contains 1 homeobox DNA-binding domain., The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation I(t(5:14)(q35;q32)) involving this gene is associated with T-cell acute lymphostatic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Isotype	IgG
Gene Name TLX3 HOX11L2 Protein Name T-cell leukemia homeobox protein 3 (Homeobox protein Hox-11L2) Immunogen Synthesized peptide derived from part region of human protein Specificity TLX3 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Reactivity	Human;Mouse
Protein Name T-cell leukemia homeobox protein 3 (Homeobox protein Hox-11L2) Immunogen Synthesized peptide derived from part region of human protein Specificity TLX3 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation (tc); 14)(q35;q32)] involving this gene is a sasciated with T-cell acute lymphoblastic leukemia (T-ALL) in children and young adults. [provided by RefSeq, Nov 2015],	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity TLX3 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation (t(5;14)(q35;q32)) involving this gene is a sasciated with T-cell acute lymphoblastic leukemia (T-ALLin children and young adults. [provided by RefSeq, Nov 2015],	Gene Name	TLX3 HOX11L2
Specificity TLX3 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5:14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALLin in children and young adults. [provided by RefSeq, Nov 2015],	Protein Name	T-cell leukemia homeobox protein 3 (Homeobox protein Hox-11L2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Specificity	TLX3 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALLin children and young adults. [provided by RefSeq, Nov 2015],	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus. Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Concentration	1 mg/ml
Synonyms Observed Band 32kD Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Purity	≥90%
Observed Band Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Synonyms	
Tissue Specificity Muscle, Function similarity:Contains 1 homeobox DNA-binding domain., The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Observed Band	32kD
Function similarity:Contains 1 homeobox DNA-binding domain., Background The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Cell Pathway	Nucleus .
The protein encoded by this gene is an orphan homeobox protein that encodes DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL in children and young adults. [provided by RefSeq, Nov 2015],	Tissue Specificity	Muscle,
in children and young adults. [provided by RefSeq, Nov 2015],	Function	similarity:Contains 1 homeobox DNA-binding domain.,
matters needing Avoid repeated freezing and thawing!	Background	The protein encoded by this gene is an orphan homeobox protein that encodes a DNA-binding nuclear transcription factor. A translocation [t(5;14)(q35;q32)] involving this gene is associated with T-cell acute lymphoblastic leukemia (T-ALL) in children and young adults. [provided by RefSeq, Nov 2015],
attention	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

