



TAF2 Polyclonal Antibody

Catalog No	BYab-06304
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	TAF2 CIF150 TAF2B
Protein Name	Transcription initiation factor TFIID subunit 2 (150 kDa cofactor of initiator function) (RNA polymerase II TBP-associated factor subunit B) (TBP-associated factor 150 kDa) (Transcription initiation f
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	TAF2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	131kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in all tissues tested.
Function	function:Transcription factor TFIID is one of the general factors required for accurate and regulated initiation by RNA polymerase II. TFIID is a multimeric protein complex that plays a central role in mediating promoter responses to various activators and repressors. It requires core promoter-specific cofactors for productive transcription stimulation. TAF2 stabilizes TFIID binding to core promoter.,miscellaneous:PubMed:9418870 was unable to show an association between TAF2 and TFIID.,sequence caution:Contaminating sequence. Potential poly-A sequence.,similarity:Belongs to the TAF2 family.,subunit:Component of transcription factor TFIID which is composed of TBP and a number of TBP-associated factors. Interacts with TAF2C1. Component of the TFTC-HAT complex.,tissue specificity:Expressed in all tissues tested.,

Nanjing BYabscience technology Co.,Ltd



Background

Initiation of transcription by RNA polymerase II requires the activities of more than 70 polypeptides. The protein that coordinates these activities is transcription factor IID (TFIID), which binds to the core promoter to position the polymerase properly, serves as the scaffold for assembly of the remainder of the transcription complex, and acts as a channel for regulatory signals. TFIID is composed of the TATA-binding protein (TBP) and a group of evolutionarily conserved proteins known as TBP-associated factors or TAFs. TAFs may participate in basal transcription, serve as coactivators, function in promoter recognition or modify general transcription factors (GTFs) to facilitate complex assembly and transcription initiation. This gene encodes one of the larger subunits of TFIID that is stably associated with the TFIID complex. It contributes to interactions at and downstream

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images