



# UIMC1 Polyclonal Antibody

<b>Catalog No</b>	BYab-06351
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	UIMC1 RAP80 RXRIP110
<b>Protein Name</b>	BRCA1-A complex subunit RAP80 (Receptor-associated protein 80) (Retinoid X receptor-interacting protein 110) (Ubiquitin interaction motif-containing protein 1)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	UIMC1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	79kD
<b>Cell Pathway</b>	Nucleus . Localizes at sites of DNA damage at double-strand breaks (DSBs).
<b>Tissue Specificity</b>	Expressed in testis, ovary, thymus and heart. Expressed in germ cells of the testis.
<b>Function</b>	domain:The Abraxas-interacting region (AIR) mediates the interaction with FAM175A/Abraxas.,domain:The UIM-linker region between the 2 UIM repeats determines the selectivity for 'Lys-63'-linked ubiquitin. The length of the linker is important. The linker reduces the flexibility between the UIM repeats and promotes high-affinity and linkage-selective interactions.,function:Ubiquitin-binding protein that specifically recognizes and binds 'Lys-63'-linked ubiquitin. Plays a central role in the BRCA1-A complex by specifically binding 'Lys-63'-linked ubiquitinated histones H2A and H2AX at DNA lesions sites, leading to target the BRCA1-BARD1 heterodimer to sites of DNA damage at double-strand breaks (DSBs). The BRCA1-A complex also possesses deubiquitinase activity that specifically removes 'Lys-63'-linked ubiquitin on

Nanjing BYabscience technology Co.,Ltd



histones H2A and H2AX. Also weakly binds monoubiquitin but with much less aff

**Background**

ubiquitin interaction motif containing 1(UIMC1) Homo sapiens This gene encodes a nuclear protein that interacts with Brca1 (breast cancer 1) in a complex to recognize and repair DNA lesions. This protein binds ubiquitinated lysine 63 of histone H2A and H2AX. This protein may also function as a repressor of transcription. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2015],

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**