



# UBA6 Polyclonal Antibody

<b>Catalog No</b>	BYab-06353
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	UBA6 MOP4 UBE1L2
<b>Protein Name</b>	Ubiquitin-like modifier-activating enzyme 6 (Ubiquitin-activating enzyme 6) (Monocyte protein 4) (MOP-4) (Ubiquitin-activating enzyme E1-like protein 2) (E1-L2)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	UBA6 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	115kD
<b>Cell Pathway</b>	cytoplasm,cytosol,integral component of membrane,
<b>Tissue Specificity</b>	Widely expressed. Isoform 2 is predominantly expressed in testis with higher expression in adult testis than in fetal testis.
<b>Function</b>	alternative products:Named isoforms=3,function:Activates ubiquitin by first adenylating its C-terminal glycine residue with ATP, and thereafter linking this residue to the side chain of a cysteine residue in E1, yielding an ubiquitin-E1 thioester and free AMP. Specific for ubiquitin, does not activate ubiquitin-like peptides. Differs from UBE1 in its specificity for substrate E2 charging. Does not charge cell cycle E2s, such as CDC34. Isoform 2 may play a key role in ubiquitin system and may influence spermatogenesis and male fertility.,pathway:Protein modification; protein ubiquitination.,similarity:Belongs to the ubiquitin-activating E1 family.,tissue specificity:Widely expressed. Isoform 2 is predominantly expressed in testis with higher expression in adult testis than in fetal testis.,

Nanjing BYabscience technology Co.,Ltd



<b>Background</b>	Modification of proteins with ubiquitin (UBB; MIM 191339) or ubiquitin-like proteins controls many signaling networks and requires a ubiquitin-activating enzyme (E1), a ubiquitin conjugating enzyme (E2), and a ubiquitin protein ligase (E3). UBE1L2 is an E1 enzyme that initiates the activation and conjugation of ubiquitin-like proteins (Jin et al., 2007 [PubMed 17597759]).[supplied by OMIM, Mar 2008],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images