



CENPI Polyclonal Antibody

Catalog No	BYab-06445
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	CENPI FSHPRH1 ICEN19 LRPR1
Protein Name	Centromere protein I (CENP-I) (FSH primary response protein 1) (Follicle-stimulating hormone primary response protein) (Interphase centromere complex protein 19) (Leucine-rich primary response protein)
Immunogen	Synthesized peptide derived from human protein . at AA range: 10-90
Specificity	CENPI Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	83kD
Cell Pathway	Nucleus. Chromosome, centromere. Localizes exclusively in the centromeres. The CENPA-CAD complex is probably recruited on centromeres by the CENPA-NAC complex.
Tissue Specificity	Epithelium,Testis,
Function	disease:May be involved in X-linked disorders of gonadal development and gametogenesis.,function:Component of the CENPA-CAD (nucleosome distal) complex, a complex recruited to centromeres which is involved in assembly of kinetochore proteins, mitotic progression and chromosome segregation. May be involved in incorporation of newly synthesized CENPA into centromeres via its interaction with the CENPA-NAC complex. Required for the localization of CENPF, MAD1L1 and MAD2 (MAD2L1 or MAD2L2) to kinetochores. Involved in the response of gonadal tissues to follicle-stimulating hormone.,induction:By follicle-stimulating hormone (FSH).,similarity:Belongs to the mis6

Nanjing BYabscience technology Co.,Ltd



family.,subcellular location:Localizes exclusively in the centromeres. The CENPA-CAD complex is probably recruited on centromeres by the CENPA-NAC complex.,subunit:Component of the CENPA-CAD complex, composed of CENPI, CENPK, CENPL, CE

Background

This gene encodes a centromere protein that is a component of the CENPA-NAC (nucleosome-associated) complex. This complex is critical for accurate chromosome alignment and segregation and it ensures proper mitotic progression. This protein regulates the recruitment of kinetochore-associated proteins that are required to generate the spindle checkpoint signal. The product of this gene is involved in the response of gonadal tissues to follicle-stimulating hormone. Mutations in this gene may be involved in human X-linked disorders of gonadal development and gametogenesis. Alternate splicing results in multiple transcript variants. A pseudogene of this gene is found on chromosome 13. [provided by RefSeq, Jan 2016],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images