




MAGAC Polyclonal Antibody

| Catalog No BYab-06471 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name MAGEA12 MAGE12 Protein Name Melanoma-associated antigen 12 (Cancer/testis antigen 1.12) (CT1.12) (MAGE-12 antigen) Immunogen Synthesized peptide derived from human protein . at AA range: 40-120 Specificity MAGAC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Synonyms Observed Band 34kD Cell Pathway Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function.Not known, though may play a role in tumor transformation or aspects of tumor progression, simi | | |
|---|---|---|
| Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name MAGEA12 MAGE12 Protein Name Melanoma-associated antigen 12 (Cancer/testis antigen 1.12) (CT1.12) (MAGE-12 antigen) (MAGE-12 anti | Catalog No | BYab-06471 |
| Applications WB;ELISA Gene Name MAGEA12 MAGE12 Protein Name (MAGE-12 antigen) (MAGE12F antigen) Immunogen Synthesized peptide derived from human protein . at AA range: 40-120 Specificity MAGAC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function Under Carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Isotype | IgG |
| Gene Name MAGEA12 MAGE12 Protein Name (MAGE-12 antigen) (MAGE12F antigen) Immunogen Synthesized peptide derived from human protein . at AA range: 40-120 Specificity MAGAC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function: Not known, though may play a role in tumor transformation or aspects of tumor progression, similarity-Contains 1 MAGE domain, tissue specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Reactivity | Human;Rat;Mouse; |
| Protein Name | Applications | WB;ELISA |
| Immunogen Synthesized peptide derived from human protein . at AA range: 40-120 Specificity MAGAC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progressionsimilarity: Contains 1 MAGE domain, tissue specificity: Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spiliced varia | Gene Name | MAGEA12 MAGE12 |
| Specificity MAGAC Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression., similarity-contains 1 MAGE domain., tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosoma X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the overexpressed in tumors. Multiple alternatively spliced variants encoding the | Protein Name | Melanoma-associated antigen 12 (Cancer/testis antigen 1.12) (CT1.12) (MAGE-12 antigen) (MAGE12F antigen) |
| Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Immunogen | Synthesized peptide derived from human protein . at AA range: 40-120 |
| Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Specificity | MAGAC Polyclonal Antibody detects endogenous levels of protein. |
| Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Source | Polyclonal, Rabbit,IgG |
| Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Purification | · |
| Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Concentration | 1 mg/ml |
| Synonyms Observed Band 34kD Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Purity | ≥90% |
| Observed Band Cell Pathway Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | | |
| Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Storage Stability | -20°C/1 year |
| Tissue Specificity Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | | -20°C/1 year |
| squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. Function function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Synonyms | |
| tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes., Background MAGE family member A12(MAGEA12) Homo sapiens This gene is closely related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Synonyms Observed Band | |
| related to several other genes clustered on chromosome X. These genes may be overexpressed in tumors. Multiple alternatively spliced variants encoding the | Synonyms Observed Band Cell Pathway | 34kD Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in |
| | Synonyms Observed Band Cell Pathway Tissue Specificity | Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but not in normal tissues except for testes. function:Not known, though may play a role in tumor transformation or aspects of tumor progression.,similarity:Contains 1 MAGE domain.,tissue specificity:Expressed in many tumors of several types, such as melanoma, head and neck squamous cell carcinoma, lung carcinoma and breast carcinoma, but |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from DU145 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658