



FOG1 Polyclonal Antibody

BYab-06524
IgG
Human;Mouse
WB;ELISA
ZFPM1 FOG1 ZFN89A
Zinc finger protein ZFPM1 (Friend of GATA protein 1) (FOG-1) (Friend of GATA 1) (Zinc finger protein 89A) (Zinc finger protein multitype 1)
Synthesized peptide derived from human protein . at AA range: 620-700
FOG1 Polyclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
1 mg/ml ≥90%
≥90%
≥90%
≥90% -20°C/1 year
≥90% -20°C/1 year 110kD

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



of some genes in gonads. May also be involved in cardiac development, in a non-redundant way with ZFPM2/FOG2.,P domain:The CCHC-type zinc fingers 1, 5, 6 and 9 directly bind to GATA-type zinc fingers. The Tyr residue adjacent to the last Cys of the CCHC-type zinc finger is Background essential for the interaction with GATA-type zinc fingers., function: Transcription regulator that plays an essential role in erythroid and megakaryocytic cell differentiation. Essential cofactor that acts via the formation of a heterodimer with transcription factors of the GATA family GATA1, GATA2 and GATA3. Such heterodimer can both activate or repress transcriptional activity, depending on the cell and promoter context. The heterodimer formed with GATA proteins is essential to activate expression of genes such as NFE2, ITGA2B, alpha- and beta-globin, while it represses expression of KLF1. May be involved in regulation of some genes in gonads. May also be involved in cardiac development, in a non-redundant way with ZFPM2/FOG2.,PTM:Phosphorylated upon DNA damage, probably by ATM or ATR.,similarity:Belongs to the FOG (Friend of GATA) family.,similarity:Contains 4 C2H2-type zinc fingers.,similarity:Contains 5 C2HC-type zinc fingers., subunit:Interacts with corepressor CTBP2; this interaction is however not essential for corepressor activity (By similarity). Interacts with the N-terminal zinc-finger of GATA1, GATA2 and probably GATA3., tissue specificity: Mainly expressed in hematopoietic tissues. Also expressed in adult cerebellum, stomach, lymph node, liver and pancreas. Expressed in fetal heart, liver and spleen., matters needing Avoid repeated freezing and thawing! attention

Usage suggestions


This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Nanjing BYabscience technology Co.,Ltd





Products Images



Western Blot analysis of Hela lysis, using primary antibody at 1:1000 dilution. Secondary antibody(catalog#:RS23920) was diluted at 1:10000

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658