



ROAA Polyclonal Antibody

Catalog No BYab-06557 Isotype IgG Reactivity Human;Mouse Applications WB;ELISA Gene Name HNRNPAB ABBP1 HNRPAB Protein Name Heterogeneous nuclear ribonucleoprotein A/B (hnRNP A/B) (APOBEC1-binding protein 1) (ABBP-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA also binds to APOBB mRNA transcripts around the RNA editing site. PTM. Dimethyl		
Reactivity Human;Mouse Applications WB;ELISA Gene Name HNRNPAB ABBP1 HNRPAB Protein Name Heterogeneous nuclear ribonucleoprotein A/B (hnRNP A/B) (APOBEC1-binding protein 1) (ABBP-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site. PTM Dimethylation at Arg-32's is probably asymmetric, sailarity Contains 2 RRM (RNA recognition motif) domains., subunit:Interacts with APOBEC1., tissue specificity: Ubiquitious. Background The RNA estimptore of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase	Catalog No	BYab-06557
Applications WB;ELISA Gene Name HNRNPAB ABBP1 HNRPAB Protein Name Heterogeneous nuclear ribonucleoprotein A/B (hnRNP A/B) (APOBEC1-binding protein 1) (ABBP-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site. PTM Dilmethylation at Arg-322 is probably asymmetric, similarity Contains 2 RRM (RNA recognition motif) domains., subunit:Interacts with APOBEC1., tissue specificity: Ubiquitous. This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear RNA (hRNA) complexes. The yere associated with pre-mRNAs in the nucleus and appear to influence pre-mRNAp rocessing and other aspects of mRNA metabolism and	Isotype	IgG
Gene Name HNRNPAB ABBP1 HNRPAB Protein Name Heterogeneous nuclear ribonucleoprotein A/B (hnRNP A/B) (APOBEC1-binding protein 1) (ABBP-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site. PTM: Dimethylation at Arg-322 is probably asymmetric, similarity: Contains 2 RRM (RNA recognition motif) domains., subunit:Interacts with APOBEC1., tissue specificityUbiquifuous, Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other assects of mRNAestolism and	Reactivity	Human;Mouse
Protein Name Heterogeneous nuclear ribonucleoprotein A/B (hnRNP A/B) (APOBEC1-binding protein 1) (ABBP-1) Immunogen Synthesized peptide derived from part region of human protein Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus. Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site., PTM: Dimethylation at Arg. 322 is probably asymmetric. similarity: Contains 2 RRM (RNA recognition motif) domains., subunit:Interacts with APOBEC1., tissue specificity:Ubiquitous. Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA metabolism and	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA . Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity. Contains 2 RRM (RNA recognition motif) domains .,subunit.Interacts with APOBEC1.,tissue specificity:Ubiquitous. Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Gene Name	HNRNPAB ABBP1 HNRPAB
Specificity ROAA Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function: Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site. PTM: Dimethylation at Arg-322 is probably asymmetric. similarity: Contains 2 RRM (RNA recognition motif) domains., subunit:Interacts with APOBEC1., tissue specificity: Ubiquitous. Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Protein Name	Heterogeneous nuclear ribonucleoprotein A/B (hnRNP A/B) (APOBEC1-binding protein 1) (ABBP-1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site. PTM: Dimethylation at Arg-322 is probably asymmetric. similarity. Contains 2 RRM (RNA recognition motif) domains., subunit:Interacts with APOBEC1., tissue specificity:Ubiquitous. Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA mRabolism and	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus Cytoplasm Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site .PTM:Dimethylation at Arg-322 is probably asymmetric .similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Inferacts with APOBEC1.,tissue specificity:Ubiquitous. Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Specificity	ROAA Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric, similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous, This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity: Ubiquitous. Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetricsimilarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site ,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Synonyms Observed Band 36kD Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Concentration	1 mg/ml
Observed Band Gell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Purity	≥90%
Observed Band Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Storage Stability	-20°C/1 year
Cell Pathway Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs. Tissue Specificity Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Synonyms	
untranslated mRNAs. Ubiquitous. Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Observed Band	36kD
Function function:Binds single-stranded RNA. Has a high affinity for G-rich and U-rich regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Cell Pathway	Nucleus . Cytoplasm . Localized in cytoplasmic mRNP granules containing untranslated mRNAs.
regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue specificity:Ubiquitous., Background This gene belongs to the subfamily of ubiquitously expressed heterogeneous nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Tissue Specificity	Ubiquitous.
nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and	Function	regions of hnRNA. Also binds to APOB mRNA transcripts around the RNA editing site.,PTM:Dimethylation at Arg-322 is probably asymmetric.,similarity:Contains 2 RRM (RNA recognition motif) domains.,subunit:Interacts with APOBEC1.,tissue
	Background	nuclear ribonucleoproteins (hnRNPs). The hnRNPs are produced by RNA polymerase II and are components of the heterogeneous nuclear RNA (hnRNA) complexes. They are associated with pre-mRNAs in the nucleus and appear to influence pre-mRNA processing and other aspects of mRNA metabolism and

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	shuttle between the nucleus and the cytoplasm. The hnRNP proteins have distinct nucleic acid binding properties. The protein encoded by this gene, which binds to one of the components of the multiprotein editosome complex, has two repeats of quasi-RRM (RNA recognition motif) domains that bind to RNAs. Two alternatively spliced transcript variants encoding different isoforms have been described for this gene. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658