



BACH2 Polyclonal Antibody

Catalog No	BYab-06565
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	BACH2
Protein Name	Transcription regulator protein BACH2 (BTB and CNC homolog 2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 250-330
Specificity	BACH2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	92kD
Cell Pathway	Cytoplasm . Nucleus . Nucleocytoplasmic shuttling is controlled by phosphorylation. .
Tissue Specificity	B-cell specific.
Function	function:Transcriptional regulator that acts as repressor or activator. Binds to Maf recognition elements (MARE). Play important roles in coordinating transcription activation and repression by MAFK.,similarity:Belongs to the bZIP family. CNC subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 1 bZIP domain.,subunit:Heterodimer of BACH2 and Maf-related transcription factors.,tissue specificity:B-cell specific.,
Background	function:Transcriptional regulator that acts as repressor or activator. Binds to Maf recognition elements (MARE). Play important roles in coordinating transcription activation and repression by MAFK.,similarity:Belongs to the bZIP family. CNC subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 1 bZIP domain.,subunit:Heterodimer of BACH2 and Maf-related transcription

Nanjing BYabscience technology Co.,Ltd



factors.,tissue specificity:B-cell specific.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images