



GBP2 Polyclonal Antibody

BYab-06611 IgG
laG
Human;Rat;Mouse;
WB;ELISA
GBP2
Interferon-induced guanylate-binding protein 2 (GTP-binding protein 2) (GBP-2) (HuGBP-2) (Guanine nucleotide-binding protein 2)
Synthesized peptide derived from human protein . at AA range: 160-240
GBP2 Polyclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
65kD
Cytoplasm . Cytoplasm, perinuclear region . Golgi apparatus membrane . Membrane; Lipid-anchor . Cytoplasmic vesicle . GBP2-GBP5 dimers localize to the Golgi apparatus
Foreskin,Lung,Skin,Spinal cord,Thymus,
function:Binds GTP, GDP and GMP. Hydrolyzes GTP very efficiently; GDP rather than GMP is the major reaction product.,induction:By interferon gamma during macrophage activation.,similarity:Belongs to the GBP family.,
This gene belongs to the guanine-binding protein (GBP) family, which includes interferon-induced proteins that can bind to guanine nucleotides (GMP, GDP and GTP). The encoded protein is a GTPase which hydrolyzes GTP, predominantly to GDP. The protein may play a role as a marker of squamous cell carcinomas. [provided by RefSeq, Jul 2013],

Nanjing BYabscience technology Co.,Ltd





matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd