




## IGFL1 Polyclonal Antibody

Catalog No         BYab-06629           Isotype         IgG           Reactivity         Human;Rat;Mouse;           Applications         WB;ELISA           Gene Name         IGFL1 UNQ644/PRO1274           Protein Name         Insulin growth factor-like family member 1           Immunogen         Synthesized peptide derived from part region of human protein AA range: 1-50           Specificity         IGFL1 Polyclonal Antibody detects endogenous levels of protein.           Formulation         Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.           Source         Polyclonal, Rabbit,IgG           Purification         The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.           Dilution         WB 1:500-2000 ELISA 1:5000-20000           Concentration         1 mg/ml           Purity         290%           Storage Stability         -20°C/1 year           Synonyms         Secreted           Cell Pathway         Secreted .           Tissue Specificity         Detected in ovary and spinal cord.           Function         similarity:Belongs to the IGFL family.           Background         IGF like family member 1(IGFL1) Homo sapiens         The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecu		
Reactivity Human;Rat;Mouse;  Applications WB;ELISA  Gene Name IGFL1 UNQ644/PRO1274  Protein Name Insulin growth factor-like family member 1  Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50  Specificity IGFL1 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  IGF like family member 1 (IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteclylically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Catalog No	BYab-06629
Applications WB;ELISA  Gene Name IGFL1 UNQ644/PRO1274  Protein Name Insulin growth factor-like family member 1  Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50  Specificity IGFL1 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.  Background IgF like family member 1 (IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor profein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Isotype	IgG
Gene Name IGFL1 UNQ644/PRO1274 Protein Name Insulin growth factor-like family member 1  Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50  Specificity IGFL1 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased	Reactivity	Human;Rat;Mouse;
Protein Name Insulin growth factor-like family member 1  Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50  Specificity IGFL1 Polyclonal Antibody detects endogenous levels of protein.  Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [Post of T cells. Increased expression of this gene may be linked to psoriasis. [Post of T cells.]	Applications	WB;ELISA
Immunogen       Synthesized peptide derived from part region of human protein AA range: 1-50         Specificity       IGFL1 Polyclonal Antibody detects endogenous levels of protein.         Formulation       Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.         Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       12kD         Cell Pathway       Secreted .         Tissue Specificity       Detected in ovary and spinal cord.         Function       similarity:Belongs to the IGFL family.,         Background       IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of 1 cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Gene Name	IGFL1 UNQ644/PRO1274
Specificity  IGFL1 Polyclonal Antibody detects endogenous levels of protein.  Formulation  Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source  Polyclonal, Rabbit, IgG  Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  12kD  Cell Pathway  Secreted.  Tissue Specificity  Detected in ovary and spinal cord.  Function  IGF like family member 1(IGFL family.,  Background  IGF like family member 1 (IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Protein Name	Insulin growth factor-like family member 1
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90% Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  Background  IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precuror protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriagis. [provided by RefSeq, Aug	Immunogen	Synthesized peptide derived from part region of human protein AA range: 1-50
Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.         Dilution       WB 1:500-2000 ELISA 1:5000-20000         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band       12kD         Cell Pathway       Secreted .         Tissue Specificity       Detected in ovary and spinal cord.         Function       similarity:Belongs to the IGFL family.,         Background       IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Specificity	IGFL1 Polyclonal Antibody detects endogenous levels of protein.
Purification  The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.  Dilution  WB 1:500-2000 ELISA 1:5000-20000  Concentration  1 mg/ml  Purity  ≥90%  Storage Stability  -20°C/1 year  Synonyms  Observed Band  12kD  Cell Pathway  Secreted .  Tissue Specificity  Detected in ovary and spinal cord.  Function  similarity:Belongs to the IGFL family.,  Background  IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen.  Dilution WB 1:500-2000 ELISA 1:5000-20000  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Source	Polyclonal, Rabbit,IgG
Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms         Observed Band       12kD         Cell Pathway       Secreted .         Tissue Specificity       Detected in ovary and spinal cord.         Function       similarity:Belongs to the IGFL family.,         Background       IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band 12kD  Cell Pathway Secreted .  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability  -20°C/1 year  Synonyms  Observed Band  12kD  Cell Pathway  Secreted.  Tissue Specificity  Detected in ovary and spinal cord.  Function  Similarity:Belongs to the IGFL family.,  Background  IGF like family member 1(IGFL1) Homo sapiens gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Concentration	1 mg/ml
Synonyms  Observed Band 12kD  Cell Pathway Secreted.  Tissue Specificity Detected in ovary and spinal cord.  Function similarity:Belongs to the IGFL family.,  Background IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Purity	≥90%
Observed Band  Cell Pathway  Secreted .  Tissue Specificity  Detected in ovary and spinal cord.  Function  Similarity:Belongs to the IGFL family.,  Background  IGF like family member 1(IGFL1) Homo sapiens gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Storage Stability	-20°C/1 year
Cell Pathway         Secreted .           Tissue Specificity         Detected in ovary and spinal cord.           Function         similarity:Belongs to the IGFL family.,           Background         IGF like family member 1(IGFL1) Homo sapiens gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Synonyms	
Tissue Specificity  Detected in ovary and spinal cord.  Function  Similarity:Belongs to the IGFL family.,  IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Observed Band	12kD
Function  Similarity:Belongs to the IGFL family.,  IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Cell Pathway	Secreted .
Background  IGF like family member 1(IGFL1) Homo sapiens The protein encoded by this gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Tissue Specificity	Detected in ovary and spinal cord.
gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug	Function	similarity:Belongs to the IGFL family.,
	Background	gene is a member of the insulin-like growth factor family of signaling molecules. The encoded protein is synthesized as a precursor protein and is proteolytically cleaved to form a secreted mature peptide. The mature peptide binds to a receptor, which in mouse was found on the cell surface of T cells. Increased expression of this gene may be linked to psoriasis. [provided by RefSeq, Aug

Nanjing BYabscience technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Nanjing BYabscience technology Co.,Ltd