



XCL2 Polyclonal Antibody

Catalog No BYab-06662 Isotype IgG Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name XCL2 SCYC2 Protein Name Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2) Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry, similarity:Belongs to the intercrine gamma family. Background dvinction:Chemotact		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name XCL2 SCYC2 Protein Name Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2) Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry, similarity:Belongs to the intercrine gamma family. Background Avoid repeated freezing and thawing!	Catalog No	BYab-06662
Applications WB;ELISA Gene Name XCL2 SCYC2 Protein Name Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2) Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry, similarity:Belongs to the intercrine gamma family., Background Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name XCL2 SCYC2 Protein Name Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2) Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood, Placenta, Function function: Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information: XCL2 entry, similarity: Belongs to the intercrine gamma family., Background function: Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information: XCL2 entry, similarity: Belongs to the intercrine gamma family., Matters needing Avoid repeated freezing and thawing!	Reactivity	Human;Rat;Mouse;
Protein Name Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2) Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils., online information:XCL2 entry, similarity:Belongs to the intercrine gamma family., Background punction:Chemotactic activity for lymphocytes but not for monocytes or neutrophils., online information:XCL2 entry, similarity:Belongs to the intercrine gamma family.	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein AA range: 1-50 Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry,similarity:Belongs to the intercrine gamma family. Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry,similarity:Belongs to the intercrine gamma family. Matters needing Avoid repeated freezing and thawing!	Gene Name	XCL2 SCYC2
Specificity XCL2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry, similarity:Belongs to the intercrine gamma family. Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils, online information:XCL2 entry, similarity:Belongs to the intercrine gamma family. matters needing Avoid repeated freezing and thawing!	Protein Name	Cytokine SCM-1 beta (C motif chemokine 2) (XC chemokine ligand 2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood, Placenta, Function function: Chemotactic activity for lymphocytes but not for monocytes or neutrophils online information: XCL2 entry, similarity: Belongs to the intercrine gamma family., Background function: Chemotactic activity for lymphocytes but not for monocytes or neutrophils. online information: XCL2 entry, similarity: Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from part region of human protein AA range: 1-50
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils., online information:XCL2 entry, similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils., online information:XCL2 entry, similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Specificity	XCL2 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry, similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry, similarity:Belongs to the intercrine gamma family., Matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 12kD Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Secreted. Tissue Specificity Blood,Placenta, Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Blood,Placenta, function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Observed Band	12kD
Function function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Cell Pathway	Secreted.
neutrophils.,online information:XĆL2 entrý,similarity:Belongs to the intercrine gamma family., Background function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family., matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Blood,Placenta,
gamma family., matters needing Avoid repeated freezing and thawing!	Function	neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine
	Background	function:Chemotactic activity for lymphocytes but not for monocytes or neutrophils.,online information:XCL2 entry,similarity:Belongs to the intercrine gamma family.,
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.



网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658