



# SYCP1 Polyclonal Antibody

<b>Catalog No</b>	BYab-06673
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Rat;Mouse;
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	SYCP1 SCP1
<b>Protein Name</b>	Synaptonemal complex protein 1 (SCP-1) (Cancer/testis antigen 8) (CT8)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	SYCP1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	107kD
<b>Cell Pathway</b>	Nucleus . Chromosome . Chromosome, centromere . In tripartite segments of synaptonemal complexes, between lateral elements in the nucleus. Its N-terminus is found towards the center of the synaptonemal complex while the C-terminus extends well into the lateral domain of the synaptonemal complex (By similarity). Only rarely detected at centromeres during leptotene and zygotene. Detected at centromeres during mid-diplotene, when it is no longer present along chromosome arms. No longer detected at centromeres at later stages of meiosis (By similarity). .
<b>Tissue Specificity</b>	Testis.
<b>Function</b>	domain:Consists of an alpha-helical stretch of 700 AA residues, flanked by N- and C-terminal globular domains. The C-terminal domain has DNA-binding capacity.,function:Major component of the transverse filaments of synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase.,subcellular location:In tripartite segments of synaptonemal complexes, between lateral elements in the nucleus. Found only where the chromosome

**Nanjing BYabscience technology Co.,Ltd**



cores are synapsed. Its N-terminus is found towards the center of the synaptonemal complex while the C-terminus extends well into the lateral domain of the synaptonemal complex.,subunit:Found in a complex with SYCE1 and SYCE2. Interacts with SYCE1 and SYCE2.,tissue specificity:Testis.,

**Background**

domain:Consists of an alpha-helical stretch of 700 AA residues, flanked by N- and C-terminal globular domains. The C-terminal domain has DNA-binding capacity.,function:Major component of the transverse filaments of synaptonemal complexes (SCS), formed between homologous chromosomes during meiotic prophase.,subcellular location:In tripartite segments of synaptonemal complexes, between lateral elements in the nucleus. Found only where the chromosome cores are synapsed. Its N-terminus is found towards the center of the synaptonemal complex while the C-terminus extends well into the lateral domain of the synaptonemal complex.,subunit:Found in a complex with SYCE1 and SYCE2. Interacts with SYCE1 and SYCE2.,tissue specificity:Testis.,

**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**