





RASEF Polyclonal Antibody

Catalog No BYab-06720 Isotype IgG Reactivity Human;Mouse Applications WB:ELISA Gene Name RASEF RAB45 Protein Name Ras and EF-hand domain-containing protein (Ras-related protein Rab-45) Immunogen Synthesized peptide derived from part region of human protein Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function GTPase superfamily. Rab family. Smillarity. Contains 2 EF-hand domains, subunit-Homodimer., issue specificity/Down-regulated in cutaneous melanoma cells but not in breast cancer cell		
Reactivity Human;Mouse Applications WB;ELISA Gene Name RASEF RAB45 Protein Name Ras and EF-hand domain-containing protein (Ras-related protein Rab-45) Immunogen Synthesized peptide derived from part region of human protein Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP, similarity:Belongs to the small GTPase superfamily. Rab family, similarity:Contains 2 EF-hand domains, subunit-Homodimer, ifsues specificity;Down-regulated in cutaneous melanoma cells but not in breast cancer cells. This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a colled-coll motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Catalog No	BYab-06720
Applications WB;ELISA Gene Name RASEF RAB45 Protein Name Ras and EF-hand domain-containing protein (Ras-related protein Rab-45) Immunogen Synthesized peptide derived from part region of human protein Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP, similarity:Belongs to the small GTPase superfamily. Rab family, similarity:Contains 2 EF-hand domains, subunit-Homodimer, tissue specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Isotype	IgG
Gene Name RASEF RAB45 Protein Name Ras and EF-hand domain-containing protein (Ras-related protein Rab-45) Immunogen Synthesized peptide derived from part region of human protein Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP, similarity:Belongs to the small GTPase superfamily. Rab family, similarity:Contains 2 EF-hand domains, subunit-Homodimer, lissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a colled-coil motif and a C-terminal Rab domanian. A	Reactivity	Human;Mouse
Protein Name Ras and EF-hand domain-containing protein (Ras-related protein Rab-45) Immunogen Synthesized peptide derived from part region of human protein Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTPsimilarity:Belongs to the small GTPase superfamily. Rab family, similarity:Contains 2 EF-hand domains, subunit-Homodimer, lissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal Er-hand domain, a colled-coil motif and a C-terminal Rab domain. A pote	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP, similarity:Belongs to the small GTPase superfamily. Rab family, similarity:Contains 2 EF-hand domains., subunit:Homodimer., tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Gene Name	RASEF RAB45
Specificity RASEF Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP., similarity:Belongs to the small GTPase superfamily. Rab family, similarity:Contains 2 EF-hand domains., subunit:Homodimer., tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a colled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq. Nov	Protein Name	Ras and EF-hand domain-containing protein (Ras-related protein Rab-45)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity \$90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function GTPase superfamily. Rab family. similarity:Contains 2 EF-hand domains, subunit:Homodimer, tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTPsimilarity:Belongs to the small GTPase superfamily. Rab familysimilarity:Contains 2 EF-hand domainssubunit.Homodimertissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Specificity	RASEF Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family,similarity:Contains 2 EF-hand domains,,subunit:Homodimer, tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region. Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Concentration	1 mg/ml
Synonyms Observed Band 81kD Cell Pathway Cytoplasm, perinuclear region. Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Purity	≥90%
Observed Band Cell Pathway Cytoplasm, perinuclear region. Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm, perinuclear region . Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., Background This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Synonyms	
Tissue Specificity Down-regulated in cutaneous melanoma cells but not in breast cancer cells. Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Observed Band	81kD
Function function:Binds predominantly GDP, and also GTP.,similarity:Belongs to the small GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Cell Pathway	Cytoplasm, perinuclear region .
GTPase superfamily. Rab family., similarity:Contains 2 EF-hand domains., subunit:Homodimer., tissue specificity:Down-regulated in cutaneous melanoma cells but not in breast cancer cells., This gene is a member of the Rab family of GTPases that are involved in regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Tissue Specificity	Down-regulated in cutaneous melanoma cells but not in breast cancer cells.
regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov	Function	GTPase superfamily. Rab family.,similarity:Contains 2 EF-hand domains.,subunit:Homodimer.,tissue specificity:Down-regulated in cutaneous
	Background	regulation of membrane traffic. The encoded protein contains an N-terminal EF-hand domain, a coiled-coil motif and a C-terminal Rab domain. A potential role as tumor suppressor has been indicated for this gene. [provided by RefSeq, Nov

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







attention	Avoid repeated freezing and triawing:
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd