



LAP2B Polyclonal Antibody

BYab-06847
lgG
Human;Mouse
WB;ELISA
TMPO LAP2
Lamina-associated polypeptide 2, isoforms beta/gamma (Thymopoietin, isoforms beta/gamma) (TP beta/gamma) (Thymopoietin-related peptide isoforms beta/gamma) (TPRP isoforms beta/gamma) [Cleaved into: Th
Synthesized peptide derived from part region of human protein
LAP2B Polyclonal Antibody detects endogenous levels of protein.
Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Polyclonal, Rabbit,IgG
The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
WB 1:500-2000 ELISA 1:5000-20000
1 mg/ml
≥90%
-20°C/1 year
49kD
Nucleus inner membrane; Single-pass type II membrane protein. Tightly associated with the nuclear lamina.; [Isoform Zeta]: Cytoplasm .
Expressed in many tissues. Most abundant in adult thymus and fetal liver.
alternative products:Additional isoforms seem to exist,domain:Has two structurally independent, non-interacting domains: LEM-like (also called LAP2-N or LEM-D) and LEM (also called LAP2-C or LEM-B). LEM-like binds DNA while LEM interacts with BANF1.,function:May be involved in the structural organization of the nucleus and in the post-mitotic nuclear assembly. Play an important role, together with LMNA, in the nuclear anchorage of RB1.,function:May help direct the assembly of the nuclear lamina and thereby help maintain the structural organization of the nuclear envelope. Possible receptor for attachment of lamin filaments to the inner nuclear membrane. May be involved in the control of initiation of DNA replication through its interaction with NAKAP95.,function:TP and

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商



BYabscience	精准的 WB 检测服务 如果
	TP5 may play a role in T-cell development and function. TP5 is an immunomodulating pentapeptide.,pharmaceutical:TP5 is a
Background	alternative products:Additional isoforms seem to exist,domain:Has two structurally independent, non-interacting domains: LEM-like (also called LAP2-N or LEM-D) and LEM (also called LAP2-C or LEM-B). LEM-like binds DNA while LEM interacts with BANF1.,function:May be involved in the structural organization of the nucleus and in the post-mitotic nuclear assembly. Play an important role, together with LMNA, in the nuclear anchorage of RB1.,function:May help direct the assembly of the nuclear lamina and thereby help maintain the structural organization of the nuclear newelope. Possible receptor for attachment of lamin filaments to the inner nuclear membrane. May be involved in the control of initiation of DNA replication through its interaction with NAKAP95.,function:TP and TP5 may play a role in T-cell development and function. TP5 is an immunomodulating pentapeptide.,pharmaceutical:TP5 is available under the names Timunox (Cilag), Sintomodulina (Italofarmaco) and Mepentil (Recordati). Used in primary and secondary immune deficiencies, autoimmunity, infections and cancer.,PTM:Mitosis-specific phosphorylation specifically abolishes its binding to lamin B and chromosomes.,PTM:Phosphorylated in a mitose-specific manner.,similarity:Contains 1 LEM-like domain.,subcellular location:Expressed diffusely throughout the nucleus.,subcellular location:Tightly associated with the nuclear lamina.,subunit:Interacts with LMNA, BANF1 and RB1 and with chromosomes. Associates directly or indirectly with lamins at specific cell-cycle stages.,subunit:Interacts with LMNB1, LMNB2, BANF1, NAKAP95, GMCL and chromosomes.,tissue specificity:Expressed in many tissues. Most abundant in adult thymus and fetal liver.,
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd