



PTPRG Polyclonal Antibody

| Catalog No | BYab-06858 |
|--------------------|--|
| Isotype | lgG |
| Reactivity | Human;Mouse |
| Applications | WB;ELISA |
| Gene Name | PTPRG PTPG |
| Protein Name | Receptor-type tyrosine-protein phosphatase gamma (Protein-tyrosine phosphatase gamma) (R-PTP-gamma) (EC 3.1.3.48) |
| Immunogen | Synthesized peptide derived from part region of human protein |
| Specificity | PTPRG Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 158kD |
| Cell Pathway | Membrane ; Single-pass type I membrane protein . |
| Tissue Specificity | Found in a variety of tissues. |
| Function | catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,similarity:Belongs to the protein-tyrosine phosphatase family. Receptor class 5 subfamily.,similarity:Contains 1 alpha-carbonic anhydrase domain.,similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 2 tyrosine-protein phosphatase domains.,tissue specificity:Found in a variety of tissues., |
| Background | The protein encoded by this gene is a member of the protein tyrosine phosphatase (PTP) family. PTPs are known to be signaling molecules that regulate a variety of cellular processes including cell growth, differentiation, mitotic cycle, and oncogenic transformation. This PTP possesses an extracellular region, a single transmembrane region, and two tandem intracytoplasmic catalytic |
| | Nanjing BYabscience technology Co.,Ltd |

Nanjing BYabscience technology Co.,Ltd

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658



国内优质抗体供应商

精准的 WB 检测服务 文义。

24H 在线服务, 欢迎咨询



| | domains, and thus represents a receptor-type PTP. The extracellular region of this PTP contains a carbonic anhydrase-like (CAH) domain, which is also found in the extracellular region of PTPRBETA/ZETA. This gene is located in a chromosomal region that is frequently deleted in renal cell carcinoma and lung carcinoma, thus is thought to be a candidate tumor suppressor gene. [provided by RefSeq, Jul 2008], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images

Nanjing BYabscience technology Co.,Ltd