



MDGA1 Polyclonal Antibody

Catalog No	BYab-06863
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	MDGA1 MAMDC3
Protein Name	MAM domain-containing glycosylphosphatidylinositol anchor protein 1 (GPI and MAM protein) (GPIM) (Glycosylphosphatidylinositol-MAM) (MAM domain-containing protein 3)
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	MDGA1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	105kD
Cell Pathway	Cell membrane ; Lipid-anchor, GPI-anchor . Associated with lipid rafts.
Tissue Specificity	Has been found in brain, heart, skeletal muscle and kidney. Found to be overexpressed in tumor tissues.
Function	function:Required for radial migration of cortical neurons in the superficial layer of the neocortex.,similarity:Contains 1 fibronectin type-III domain.,similarity:Contains 1 MAM domain.,similarity:Contains 6 Ig-like (immunoglobulin-like) domains.,subcellular location:Associated with lipid rafts.,subunit:Interacts heterophilically through its MAM domain with proteins in axon-rich regions and through its Ig-like domains with proteins in differentiating muscle.,tissue specificity:Has been found in brain, heart, skeletal muscle and kidney. Found to be overexpressed in tumor tissues.,
Background	This gene encodes a glycosylphosphatidylinositol (GPI)-anchored cell surface glycoprotein that is expressed predominantly in the developing nervous system. In
	N " BY I ' (I I G I I I

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	addition to possessing several cell adhesion molecule-like domains, the mature protein has six Ig-like domains, a single fibronectin type III domain, a MAM domain and a C-terminal GPI-anchoring site. Studies in other mammals suggest this protein plays a role in cell adhesion, migration, and axon guidance and, in the developing brain, neuronal migration. In humans, this gene is associated with bipolar disorder and schizophrenia. [provided by RefSeq, Oct 2016],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd