



NEUS Polyclonal Antibody

neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or		
Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name SERPINI1 PI12 Protein Name Neuroserpin (Peptidase inhibitor 12) (PI-12) (Serpin I1) Immunogen Synthesized peptide derived from part region of human protein Specificity NEUS Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed: 17040209). Predominantly expressed in the brain (PubMed: 9070919). Function disease: Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN18) [MIM:604218]. FEN18 is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin. function: Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adurerous glasmining and activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adurerous glasmining and activator, similarity. Belongs to the serpin family, tissue	Catalog No	BYab-06885
Applications Gene Name SERPINI1 PI12 Protein Name Neuroserpin (Peptidase inhibitor 12) (PI-12) (Serpin I1) Immunogen Synthesized peptide derived from part region of human protein Specificity NEUS Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neurosepin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin. function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the aduneryous system. May protect neurons from cell damage by tissue—type plasminogen activators. similarly: Belongs to the serpin famility, tissue	Isotype	IgG
Gene Name SERPINI1 PI12 Protein Name Neuroserpin (Peptidase inhibitor 12) (PI-12) (Serpin I1) Immunogen Synthesized peptide derived from part region of human protein Specificity NEUS Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPIN11 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin. function:Serine protections as well as for synaptic oplasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the aduneryous system. May protect neurons from cell damage by tissue-type plasminogen activators. Serine protect neurons from cell damage by tissue-type	Reactivity	Human;Rat;Mouse
Protein Name Neuroserpin (Peptidase inhibitor 12) (PI-12) (Serpin I1) Immunogen Synthesized peptide derived from part region of human protein Specificity NEUS Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN18) [Mill.604218]. FEN18 is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies (FEN18) [Mill.604218]. FEN18 is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies on biochemically by polymers of neuroserpin., function: Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adunerous system. May protect neurons from cell damage by tissue-type plasminogen activators, similarity:Belongs to the serpin family, tissue	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity NEUS Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies and biochemically by polymers of neuroserpin inclusion bodies and biochemically by polymers of neuroserpin inclusion bodies and biochemically by polymers of neuroserpin family incution. Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adurency of synaptic connections as well as for synaptic plasticity in	Gene Name	SERPINI1 PI12
Specificity NEUS Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by opimers of neuroserpin., function: Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adureryous system. May protect neurons from cell damage by tissue-type plasminogen activators. Similiarity:Belongs to the serpin family., tissue	Protein Name	Neuroserpin (Peptidase inhibitor 12) (PI-12) (Serpin I1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted. Cytoplasmic vesicle, secretory vesicle lumen. Perikaryon. Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPIN11 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin., function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adureryous system. May protect neurons from cell damage by tissue-type plasminogen activators, similarity:Belongs to the serpin familly, tissue	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuroserpin. function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the aduneryous system. May protect neurons from cell damage by tissue-type plasminogen activator, similarity:Belongs to the serpin family, tissue	Specificity	NEUS Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin. function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adunervous system. May protect neurons from cell damage by tissue-type plasminogen activators, similarity: Belongs to the serpin family., tissue	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin., function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adunervous system. May protect neurons from cell damage by tissue-type plasminogen activator., similarity:Belongs to the serpin familly, tissue	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin., function: Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adu nervous system. May protect neurons from cell damage by tissue-type plasminogen activator., similarity: Belongs to the serpin family, tissue	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin., function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adu nervous system. May protect neurons from cell damage by tissue-type plasminogen activator., similarity:Belongs to the serpin family., tissue	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin., function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adureryous system. May protect neurons from cell damage by tissue-type plasminogen activator:, similarity:Belongs to the serpin family, tissue	Concentration	1 mg/ml
Synonyms Observed Band 45kD Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adu nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue	Purity	≥90%
Observed Band Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adu nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue	Storage Stability	-20°C/1 year
Cell Pathway Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon . Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adult nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue	Synonyms	
Tissue Specificity Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Gisease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adule nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue	Observed Band	45kD
(PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919). Function disease:Defects in SERPINI1 are the cause of familial encephalopathy with neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adu nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue	Cell Pathway	Secreted . Cytoplasmic vesicle, secretory vesicle lumen . Perikaryon .
neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adu nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue	Tissue Specificity	Detected in brain cortex and hippocampus pyramidal neurons (at protein level) (PubMed:17040209). Predominantly expressed in the brain (PubMed:9070919).
	Function	neuroserpin inclusion bodies (FEN1B) [MIM:604218]. FEN1B is characterized clinically as an autosomal dominantly inherited dementia, histologically by unique neuronal inclusion bodies and biochemically by polymers of neuroserpin.,function:Serine protease inhibitor that inhibits plasminogen activators and plasmin but not thrombin. May be involved in the formation or reorganization of synaptic connections as well as for synaptic plasticity in the adult nervous system. May protect neurons from cell damage by tissue-type plasminogen activator.,similarity:Belongs to the serpin family.,tissue

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	This gene encodes a member of the serpin superfamily of serine proteinase inhibitors. The protein is primarily secreted by axons in the brain, and preferentially reacts with and inhibits tissue-type plasminogen activator. It is thought to play a role in the regulation of axonal growth and the development of synaptic plasticity. Mutations in this gene result in familial encephalopathy with neuroserpin inclusion bodies (FENIB), which is a dominantly inherited form of familial encephalopathy and epilepsy characterized by the accumulation of mutant neuroserpin polymers. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658