



# TP4A1 Polyclonal Antibody

<b>Catalog No</b>	BYab-06967
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse;Rat
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	PTP4A1 PRL1 PTPCAAX1
<b>Protein Name</b>	Protein tyrosine phosphatase type IVA 1 (EC 3.1.3.48) (PTP(CAAXI)) (Protein-tyrosine phosphatase 4a1) (Protein-tyrosine phosphatase of regenerating liver 1) (PRL-1)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	TP4A1 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	19kD
<b>Cell Pathway</b>	Cell membrane ; Lipid-anchor . Early endosome . Endoplasmic reticulum . Cytoplasm . Cytoplasm, cytoskeleton, spindle . Nucleus . And mitotic spindle. .
<b>Tissue Specificity</b>	Expressed in bone marrow, lymph nodes, T lymphocytes, spleen, thymus and tonsil. Overexpressed in tumor cell lines.
<b>Function</b>	catalytic activity:Protein tyrosine phosphate + H(2)O = protein tyrosine + phosphate.,developmental stage:Expressed in fetal liver.,enzyme regulation:Inhibited by sodium orthovanadate and pentamidine.,function:Protein tyrosine phosphatase which stimulates progression from G1 into S phase during mitosis. May play a role in the development and maintenance of differentiating epithelial tissues. Enhances cell proliferation, cell motility and invasive activity, and promotes cancer metastasis.,induction:Strongly down-regulated upon tetrodotoxin treatment.,PTM:Farnesylated. Farnesylation is required for membrane targeting. Unfarnesylated forms are shifted into the nucleus.,similarity:Belongs to the protein-tyrosine phosphatase


Nanjing BYabscience technology Co.,Ltd




	family.,similarity:Contains 1 tyrosine-protein phosphatase domain.,subcellular location:And mitotic spindle.,subunit:Homotrimer. Interacts with ATF5 (By similarity). Inte
<b>Background</b>	protein tyrosine phosphatase type IVA, member 1(PTP4A1) Homo sapiens This gene encodes a member of a small class of prenylated protein tyrosine phosphatases (PTPs), which contain a PTP domain and a characteristic C-terminal prenylation motif. The encoded protein is a cell signaling molecule that plays regulatory roles in a variety of cellular processes, including cell proliferation and migration. The protein may also be involved in cancer development and metastasis. This tyrosine phosphatase is a nuclear protein, but may associate with plasma membrane by means of its prenylation motif. Pseudogenes related to this gene are located on chromosomes 1, 2, 5, 7, 11 and X. [provided by RefSeq, Jun 2013],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.




## Products Images



Li, X., Ma, N., Zhang, Y. et al. Circular RNA circNRIP1 promotes migration and invasion in cervical cancer by sponging miR-629-3p and regulating the PTP4A1/ERK1/2 pathway. *Cell Death Dis* 11, 399 (2020).



Li, X., Ma, N., Zhang, Y. et al. Circular RNA circNRIP1 promotes migration and invasion in cervical cancer by sponging miR-629-3p and regulating the PTP4A1/ERK1/2 pathway. *Cell Death Dis* 11, 399 (2020).



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night