





EDN2 Polyclonal Antibody

Catalog No BYab-07002 Isotype IgG Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name EDN2 Protein Name Endothelin-2 (ET-2) (Preproendothelin-2) (PPET2) Immunogen Synthesized peptide derived from part region of human protein Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Coll Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function: Endothelins are endothelium-derived vasoconstrictor peptides. similarity Belongs to the endothelin/searafotoxin family, tissue specificity/ Expressed in lung, but not in placental stem vil		
Reactivity Human;Rat;Mouse Applications WB;ELISA Gene Name EDN2 Protein Name Endothelin-2 (ET-2) (Preproendothelin-2) (PPET2) Immunogen Synthesized peptide derived from part region of human protein Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Puriffication The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villismooth muscle cells. Function function.Endothelins are endothelium-derived vasoconstrictor peptides, similarity/Belongs to the endothelin/sarafotoxin family, tissue specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villismooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigipressis. Alternative splicing results in multiple transcript	Catalog No	BYab-07002
Applications WB;ELISA Gene Name EDN2 Protein Name Endothelin-2 (ET-2) (Preproendothelin-2) (PPET2) Immunogen Synthesized peptide derived from part region of human protein Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villismooth muscle cells. Function function-Endothelins are endothelium-derived vasoconstrictor peptides, similarity/Belongs to the endothelin/sarafotoxin family, tissue specificity (Expressed in lung, but not in placental stem villi vessels or cultured placental villismooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigipressis. Alternative splicing results in multiple transcript	Isotype	IgG
Protein Name EDN2 Protein Name Endothelin-2 (ET-2) (Preproendothelin-2) (PPET2) Immunogen Synthesized peptide derived from part region of human protein Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides., similarity:Belongs to the endothelin/sarafotoxin family, tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript.	Reactivity	Human;Rat;Mouse
Protein Name Endothelin-2 (ET-2) (Preproendothelin-2) (PPET2) Immunogen Synthesized peptide derived from part region of human protein Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides, similarity. Belongs to the endothelin/sarafotoxin family, tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signalling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Applications	WB;ELISA
Immunogen Synthesized peptide derived from part region of human protein Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides, similarity Belongs to the endothelin/sarafotoxin family, tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin receptors that intale intracellular signaling events. This gene product is involved in a wider angle of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Gene Name	EDN2
Specificity EDN2 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function: Endothelins are endothelium-derived vasoconstrictor peptidessimilarity:Belongs to the endothelin/sarafotoxin familytissue specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Protein Name	Endothelin-2 (ET-2) (Preproendothelin-2) (PPET2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptidessimilarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript.	Immunogen	Synthesized peptide derived from part region of human protein
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides, similarity:Belongs to the endothelin/sarafotoxin family, tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Specificity	EDN2 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptidessimilarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Concentration	1 mg/ml
Synonyms Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function: Endothelins are endothelium-derived vasoconstrictor peptides., similarity: Belongs to the endothelin/sarafotoxin family., tissue specificity: Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Purity	≥90%
Observed Band 19kD Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Storage Stability	-20°C/1 year
Cell Pathway Secreted. Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Synonyms	
Tissue Specificity Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells. Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Observed Band	19kD
Function function:Endothelins are endothelium-derived vasoconstrictor peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Cell Pathway	Secreted.
peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured placental villi smooth muscle cells., Background This gene encodes a member of the endothelin protein family of secretory vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Tissue Specificity	
vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript	Function	peptides.,similarity:Belongs to the endothelin/sarafotoxin family.,tissue specificity:Expressed in lung, but not in placental stem villi vessels or cultured
	Background	vasoconstrictive peptides. The preproprotein is processed to a short mature form which functions as a ligand for the endothelin receptors that initiate intracellular signaling events. This gene product is involved in a wide range of biological processes, such as hypertension and ovulation. Altered expression of this gene is implicated in tumorigenesis. Alternative splicing results in multiple transcript

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658






matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from KB cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd