



BMPER Polyclonal Antibody

Catalog No BYab-07103 Isotype IgG Reactivity Human;Mouse Applications WB:ELISA Gene Name BMPER KIAA1965 Protein Name BMP-binding endothelial regulator protein (Bone morphogenetic protein-binding endothelial cell precursor-derived regulator) (Protein crossveinless-2) (hCV2) Immunogen Synthesized peptide derived from human protein . at AA range: 440-520 Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted BMP responsiveness of osteoblasts and chondrocytes, similarity:Contains 1 Tilladional, it may regulated the protein in the hondrogenic cells. Multations in this gene are associated with a lethal skeleta disorder, diaphanespon		
Reactivity Human;Mouse Applications WB;ELISA Gene Name BMPER KIAA1965 Protein Name BMP-binding endothelial regulator protein (Bone morphogenetic protein-binding endothelial cell precursor-derived regulator) (Protein crossveinless-2) (hCV2) Immunogen Synthesized peptide derived from human protein . at AA range: 440-520 Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes. similarity:Contains 1 TIL (trypsin inhibitory-like) domain, similarity:Contains 1 WFD domain similarity:Contains 1 WF	Catalog No	BYab-07103
Applications Gene Name BMPER KIAA1965 Protein Name BMP-binding endothelial regulator protein (Bone morphogenetic protein-binding endothelial cell precursor-derived regulator) (Protein crossveinless-2) (hCV2) Immunogen Synthesized peptide derived from human protein at AA range: 440-520 Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted. Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function.Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes. similarity:Contains 1 TIL (trypsin inhibitory-like) domain, similarity:Contains 1 WWFD domain, similarity:Contains 1	Isotype	IgG
Gene Name BMPER KIAA1965 Protein Name BMP-binding endothelial regulator protein (Bone morphogenetic protein-binding endothelial cell precursor-derived regulator) (Protein crossveinless-2) (hCv2) Immunogen Synthesized peptide derived from human protein . at AA range: 440-520 Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes, similarity:Contains 1 **WPD domain, similarity:Conta	Reactivity	Human;Mouse
Protein Name BMP-binding endothelial regulator protein (Bone morphogenetic protein-binding endothelial cell precursor-derived regulator) (Protein crossveinless-2) (hCV2) Immunogen Synthesized peptide derived from human protein . at AA range: 440-520 Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes, similarity:Contains 1 TIL (trypsin inhibitory-like) domain, similarity:Contains 1 WWFD domain, similarity:Contains 5 WWFC domains, subunit:Interacts with BMP4, tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes. Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiations in this gene are associated with a lethal skelefate the second of the chondrogenic cells. Mutations in this gene are associated with a lethal skelefate the second of the chondrogenic cells. Mutations in this gene are associated with a lethal skelefate the second of the chondrogenic cells. Mutations in this gene are associated with a lethal skelefate the second of the chondrogenic cells. Mutations in this gene are associated with a lethal skelefate the second of the chondrogenic cells. Mutations in this gene are associated with a lethal skelefate the second of the chondrogenic cell	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 440-520 Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Secreted . Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes, similarity:Contains 1 TIL (trypsin inhibitory-like) domain, similarity:Contains 1 TWFD domain, similarity:Contains 1 TWFD domain, similarity:Contains 3 TWFC domains, subunit:Interacts with BMP4., issue specificity. Highly expressed in lung, and brain and also in primary chondrocytes., imilarity: cells: Mutations in this gene are associated with a lethal skeletate of the chondrogenic cells. Mutations in this gene are associated with a lethal skeletate of the chondrogenic cells.	Gene Name	BMPER KIAA1965
Specificity BMPER Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes, similarity:Contains 1 TIL (trypsin inhibitory-like) domain, similarity:Contains 5 VWFC domains, subunit:Interacts with BMP4-, tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes. This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate when the colondrogenic cells. Mutations in this gene are associated with a lethal skeletate.	Protein Name	BMP-binding endothelial regulator protein (Bone morphogenetic protein-binding endothelial cell precursor-derived regulator) (Protein crossveinless-2) (hCV2)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted. Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes, similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 5 VWFC domains, subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes. Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent of the chondrogenic cells. Mutations in this gene a associated with a lethal skeleta	Immunogen	Synthesized peptide derived from human protein . at AA range: 440-520
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 1 TWFD domain., similarity:Contains 5 VWFC domains, subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes. Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP4-dependent differentiation of the chondrogenic cells. Whutations in this gene are associated with a lethal skeletated with a lethal skeletated with a lethal skeletated with a lethal skeletated with a lethal skeletated.	Specificity	BMPER Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes.,similarity:Contains 1 TIL (trypsin inhibitory-like) domain.,similarity:Contains 1 VWFD domain.,similarity:Contains 5 VWFC domains.,subunit:Interacts with BMP4.,tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted. Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 1 VWFD domain., similarity:Contains 5 VWFC domains, subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 5 VWFC domains., subunit:Interacts with BMP4.itssue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes.,similarity:Contains 1 TIL (trypsin inhibitory-like) domain.,similarity:Contains 1 VWFD domain.,similarity:Contains 5 VWFC domains.,subunit:Interacts with BMP4.,tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 75kD Cell Pathway Secreted. Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 1 VWFD domain., similarity:Contains 5 VWFC domains., subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Concentration	1 mg/ml
Synonyms Observed Band 75kD Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 1 VWFD domain., similarity:Contains 5 VWFC domains., subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Purity	≥90%
Observed Band Cell Pathway Secreted . Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes.,similarity:Contains 1 TIL (trypsin inhibitory-like) domain.,similarity:Contains 1 VWFD domain.,similarity:Contains 5 VWFC domains.,subunit:Interacts with BMP4.,tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Storage Stability	-20°C/1 year
Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 1 VWFD domain., similarity:Contains 5 VWFC domains., subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Synonyms	
Tissue Specificity Highly expressed in lung, and brain and also in primary chondrocytes. function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes.,similarity:Contains 1 TIL (trypsin inhibitory-like) domain.,similarity:Contains 1 VWFD domain.,similarity:Contains 5 VWFC domains.,subunit:Interacts with BMP4.,tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Observed Band	75kD
Function function:Inhibitor of bone morphogenetic protein (BMP) function, it may regulate BMP responsiveness of osteoblasts and chondrocytes.,similarity:Contains 1 TIL (trypsin inhibitory-like) domain.,similarity:Contains 1 VWFD domain.,similarity:Contains 5 VWFC domains.,subunit:Interacts with BMP4.,tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Cell Pathway	Secreted .
BMP responsiveness of osteoblasts and chondrocytes., similarity:Contains 1 TIL (trypsin inhibitory-like) domain., similarity:Contains 1 VWFD domain., similarity:Contains 5 VWFC domains., subunit:Interacts with BMP4., tissue specificity:Highly expressed in lung, and brain and also in primary chondrocytes., Background This gene encodes a secreted protein that interacts with, and inhibits bone morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Tissue Specificity	Highly expressed in lung, and brain and also in primary chondrocytes.
morphogenetic protein (BMP) function. It has been shown to inhibit BMP2- and BMP4-dependent osteoblast differentiation and BMP-dependent differentiation of the chondrogenic cells. Mutations in this gene are associated with a lethal skeleta	Function	BMP responsiveness of osteoblasts and chondrocytes.,similarity:Contains 1 TIL (trypsin inhibitory-like) domain.,similarity:Contains 1 VWFD domain.,similarity:Contains 5 VWFC domains.,subunit:Interacts with BMP4.,tissue specificity:Highly expressed in lung, and brain and also in primary
	Background	

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







attention	Avoid repeated freezing and triawing:
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd