




OTUD4 Polyclonal Antibody

Catalog No BYab-07111 Isotype IgG Reactivity Human;Mouse Applications WB;ELISA Gene Name OTUD4 HIN1 KIAA1046 Protein Name OTU domain-containing protein 4 (HIV-1-induced protein HIN-1) Immunogen Synthesized peptide derived from human protein . at AA range: 910-990 Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, elemention in a chimeric transcript in HIV-1 infected cells. The premature stop may be due to intron retention . sequence caution: Contaminating sequence. Originates from a chimeric transcript variants have been found for t		
Reactivity Human;Mouse Applications WB;ELISA Gene Name OTUD4 HIN1 KIAA1046 Protein Name OTU domain-containing protein 4 (HIV-1-induced protein HIN-1) Immunogen Synthesized peptide derived from human protein . at AA range: 910-990 Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit.lgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, induction: By HIV-1 insertion, sequence caution: Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention Sequence caution: Intron retention., sequence cauti	Catalog No	BYab-07111
Applications WB;ELISA Gene Name OTUD4 HIN1 KIAA1046 Protein Name OTU domain-containing protein 4 (HIV-1-induced protein HIN-1) Immunogen Synthesized peptide derived from human protein . at AA range: 910-990 Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention . sequence caution: Contaminating sequence Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention . sequence caution: Contaminating sequence Cortention is 1 of U domain. Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010],	Isotype	IgG
Gene Name OTUD4 HIN1 KIAA1046 Protein Name OTU domain-containing protein 4 (HIV-1-induced protein HIN-1) Immunogen Synthesized peptide derived from human protein . at AA range: 910-990 Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, induction: By HIV-1 insertion ., sequence caution: Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention ., sequence caution: Contaminating sequence. Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010],	Reactivity	Human;Mouse
Protein Name OTU domain-containing protein 4 (HIV-1-induced protein HIN-1) Immunogen Synthesized peptide derived from human protein . at AA range: 910-990 Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 of UTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010],	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 910-990 Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, Function induction:By HIV-1 insertion . sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention, sequence caution:Intron retention, similarity:Contains 1 OTU domain. Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010],	Gene Name	OTUD4 HIN1 KIAA1046
Specificity OTUD4 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, Function induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention, sequence caution:Intron retention.,similarity:Contains 1 OTU domain. Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [Provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Protein Name	OTU domain-containing protein 4 (HIV-1-induced protein HIN-1)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, induction:By HIV-1 insertion., sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention., sequence caution:Intron retention., similarity:Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human protein . at AA range: 910-990
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, Function induction:By HIV-1 insertion .,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention sequence caution:Intron retentionsimilarity:Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Specificity	OTUD4 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic. Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic . Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, Function induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms -20°C/1 year Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, Function induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], Matters needing Avoid repeated freezing and thawing!	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms Observed Band 122kD Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, induction: By HIV-1 insertion., sequence caution: Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention., sequence caution: Intron retention., similarity: Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Tissue Specificity Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue, induction: By HIV-1 insertion., sequence caution: Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention., sequence caution: Intron retention., similarity: Contains 1 OTU domain., Background Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Nucleus . Primarily cytoplasmic Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], Matters needing Avoid repeated freezing and thawing!	Synonyms	
Tissue Specificity Brain,Colon carcinoma,Epithelium,Mammary carcinoma,Testis,Tongue, induction:By HIV-1 insertion.,sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention.,sequence caution:Intron retention.,similarity:Contains 1 OTU domain., Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], Matters needing Avoid repeated freezing and thawing!	Observed Band	122kD
Function induction:By HIV-1 insertion., sequence caution:Contaminating sequence. Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention., sequence caution:Intron retention., similarity:Contains 1 OTU domain., Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], Matters needing Avoid repeated freezing and thawing!	Cell Pathway	Cytoplasm . Nucleus . Primarily cytoplasmic
Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention., sequence caution: Intron retention., similarity: Contains 1 OTU domain., Alternatively spliced transcript variants have been found for this gene. The smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Tissue Specificity	Brain, Colon carcinoma, Epithelium, Mammary carcinoma, Testis, Tongue,
smaller protein isoform encoded by the shorter transcript variant is found only in HIV-1 infected cells. [provided by RefSeq, Jul 2010], matters needing Avoid repeated freezing and thawing!	Function	Originates from a chimeric transcript in HIV-1-infected cells. The premature stop may be due to intron retention., sequence caution: Intron
	Background	smaller protein isoform encoded by the shorter transcript variant is found only in
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd







Usage suggestio	ns
-----------------	----

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

	Produc	ts Images		

Nanjing BYabscience technology Co.,Ltd