



HGFL Polyclonal Antibody

Catalog No	BYab-07164
lsotype	lgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	MST1 D3F15S2 DNF15S2 HGFL
Protein Name	Hepatocyte growth factor-like protein (Macrophage stimulatory protein) (Macrophage-stimulating protein) (MSP) [Cleaved into: Hepatocyte growth factor-like protein alpha chain; Hepatocyte growth factor
Immunogen	Synthesized peptide derived from human protein . at AA range: 420-500
Specificity	HGFL Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Purity Storage Stability	≥90% -20°C/1 year
-	
Storage Stability	
Storage Stability Synonyms	-20°C/1 year
Storage Stability Synonyms Observed Band	-20°C/1 year 78kD
Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year 78kD Secreted.
Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year 78kD Secreted. Liver,Plasma,Testis, function:Probably has no proteolytic activity, since crucial characteristic of serine proteases catalytic sites are not conserved.,PTM:May be cleaved after Arg-483, to yield two chains held together by disulfide bonds, or two separate polypeptides.,similarity:Belongs to the peptidase S1 family.,similarity:Belongs to the peptidase S1 family. Plasminogen subfamily.,similarity:Contains 1 PAN domain.,similarity:Contains 1 peptidase S1 domain.,similarity:Contains 4 kringle domains.,subunit:Dimer of an alpha chain and a beta chain linked by a disulfide

网址:www.njbybio.com 官方热线:025-5229-8998 监督电话:15950492658



国内优质抗体供应商

精准的 WB 检测服务

24H 在线服务,欢迎咨询



	presence of the serine protease domain, the encoded protein may not have any proteolytic activity. The receptor for this protein is RON tyrosine kinase, which upon activation stimulates ciliary motility of ciliated epithelial lung cells. This protein is secreted and cleaved to form an alpha chain and a beta chain bridged by disulfide bonds. [provided by RefSeq, Jan 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd