



AMGO1 Polyclonal Antibody

Catalog No	BYab-07233
Isotype	IgG
Reactivity	Human;Rat;Mouse
Applications	WB;ELISA
Gene Name	AMIGO1 ALI2 AMIGO KIAA1163
Protein Name	Amphoterin-induced protein 1 (AMIGO-1) (Alivin-2)
Immunogen	Synthesized peptide derived from human protein . at AA range: 120-200
Specificity	AMGO1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	54kD
Cell Pathway	Cell membrane ; Single-pass type I membrane protein . Perikaryon . Cell projection, dendrite . Cell projection, axon . Colocalizes with KCNB1 at high-density somatodendritic clusters on the surface of hippocampal and cortical neurons. Associated with axons of neuronal cells. .
Tissue Specificity	Brain,Hippocampus,Monocytic leukemia,Ovary,
Function	function:Promotes growth and fasciculation of neurites from cultured hippocampal neurons. May be involved in fasciculation as well as myelination of developing neural axons. May have a role in regeneration as well as neural plasticity in the adult nervous system. May mediate homophilic as well as heterophilic cell-cell interaction and contribute to signal transduction through its intracellular domain.,similarity:Belongs to the immunoglobulin superfamily. AMIGO family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 6 LRR (leucine-rich) repeats.,subcellular location:Associated with axons of neuronal cells.,subunit:Binds itself as well as AMIGO2 and AMIGO3.,

Nanjing BYabscience technology Co.,Ltd



Background

function:Promotes growth and fasciculation of neurites from cultured hippocampal neurons. May be involved in fasciculation as well as myelination of developing neural axons. May have a role in regeneration as well as neural plasticity in the adult nervous system. May mediate homophilic as well as heterophilic cell-cell interaction and contribute to signal transduction through its intracellular domain.,similarity:Belongs to the immunoglobulin superfamily. AMIGO family.,similarity:Contains 1 Ig-like C2-type (immunoglobulin-like) domain.,similarity:Contains 6 LRR (leucine-rich) repeats.,subcellular location:Associated with axons of neuronal cells.,subunit:Binds itself as well as AMIGO2 and AMIGO3.,

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images