



# HMGA2 Polyclonal Antibody

<b>Catalog No</b>	BYab-07296
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	HMGA2 HMGIC
<b>Protein Name</b>	High mobility group protein HMGI-C (High mobility group AT-hook protein 2)
<b>Immunogen</b>	Synthesized peptide derived from human protein . at AA range: 11-60
<b>Specificity</b>	HMGA2 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	11kD
<b>Cell Pathway</b>	Nucleus.
<b>Tissue Specificity</b>	Aorta endothelial cell,Hepatoma,
<b>Function</b>	developmental stage:Expressed predominantly during embryogenesis.,disease:A chromosomal aberration involving HMGA2 is associated with a subclass of benign mesenchymal tumors known as lipomas. Translocation t(3;12)(q27-q28;q13-q15) with LPP is shown in lipomas. HMGA2 is also fused with a number of other genes in lipomas.,disease:A chromosomal aberration involving HMGA2 is associated with parosteal lipomas. Translocation t(3;12)(q28;q14) with LPP is also shown in one parosteal lipoma.,disease:A chromosomal aberration involving HMGA2 is associated with pulmonary chondroid hamartomas. Translocation t(3;12)(q27-q28;q14-q15) with LPP is detected in pulmonary chondroid hamartomas.,disease:A chromosomal aberration involving HMGA2 is found in uterine leiomyoma (UL) [MIM:150699]. Translocation t(12;14)(q15;q23-24) with RAD51L1. Chromosomal rearrangements involving HMGA2 do not seem to be

Nanjing BYabscience technology Co.,Ltd



the princ

**Background**

This gene encodes a protein that belongs to the non-histone chromosomal high mobility group (HMG) protein family. HMG proteins function as architectural factors and are essential components of the enhancosome. This protein contains structural DNA-binding domains and may act as a transcriptional regulating factor. Identification of the deletion, amplification, and rearrangement of this gene that are associated with myxoid liposarcoma suggests a role in adipogenesis and mesenchymal differentiation. A gene knock out study of the mouse counterpart demonstrated that this gene is involved in diet-induced obesity. Alternate transcriptional splice variants, encoding different isoforms, have been characterized. [provided by RefSeq, Jul 2008],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from HCT116 cells, primary antibody was diluted at 1:1000, 4° over night