




OR4D9 Polyclonal Antibody

family., Olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory rece proteins are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Isotype IgG		
Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name OR4D9 Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamilly., olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptore proteins are members of a large family of G-protein-coupled receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors are members of a large family of G-protein-coupled receptors and are responsible for the recognition and G protein-mediated	Reactivity Human;Rat;Mouse; Applications WB;ELISA Gene Name OR4D9 Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Isotype	BYab-07482
Applications WB;ELISA Gene Name OR4D9 Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPC arising from single coding-exon genes. Olfactory receptors Share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Applications WB;ELISA Gene Name OR4D9 Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	7 .	IgG
Gene Name OR4D9 Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptorsimilarity:Belongs to the G-protein coupled recefamilly., Background Olfactory receptor family 4 subfamily D member 9(OR4D9) Home sapiens olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors interact with many neurotransmitter and hormor receptors and are	Gene NameOR4D9Protein NameOlfactory receptor 4D9 (Olfactory receptor OR11-253)ImmunogenSynthesized peptide derived from human protein . at AA range: 170-250SpecificityOR4D9 Polyclonal Antibody detects endogenous levels of protein.FormulationLiquid in PBS containing 50% glycerol, and 0.02% sodium azide.SourcePolyclonal, Rabbit, IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsSynonymsObserved Band34kDCell PathwayCell membrane; Multi-pass membrane protein.	Reactivity	Human;Rat;Mouse;
Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptarily., olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens olfactory receptors receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors (GPC arising from single coding-exon genes. Olfactory receptors share a responsible for the recognition and proterior and hormor receptors and are responsible for the recognition and proterior and hormor receptors and are responsible	Protein Name Olfactory receptor 4D9 (Olfactory receptor OR11-253) Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Applications	WB;ELISA
Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor family., olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a protein-co	Immunogen Synthesized peptide derived from human protein . at AA range: 170-250 Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Gene Name	OR4D9
Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Specificity OR4D9 Polyclonal Antibody detects endogenous levels of protein. Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Protein Name	Olfactory receptor 4D9 (Olfactory receptor OR11-253)
Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptamily., olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors of a large family of G-protein-coupled receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Formulation Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Immunogen	Synthesized peptide derived from human protein . at AA range: 170-250
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptories are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.DilutionWB 1:500-2000 ELISA 1:5000-20000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsObserved Band34kDCell PathwayCell membrane; Multi-pass membrane protein.	Specificity	OR4D9 Polyclonal Antibody detects endogenous levels of protein.
Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptores in a members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Purification The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein.	Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors Share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	affinity-chromatography using epitope-specific immunogen. Dilution WB 1:500-2000 ELISA 1:5000-20000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein.	Purification	·
Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein.	Dilution	WB 1:500-2000 ELISA 1:5000-20000
Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptore in a remembers of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Storage Stability -20°C/1 year Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Concentration	1 mg/ml
Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Synonyms Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Purity	≥90%
Observed Band Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background Olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Observed Band 34kD Cell Pathway Cell membrane; Multi-pass membrane protein.	Storage Stability	-20°C/1 year
Cell Pathway Cell membrane; Multi-pass membrane protein. Tissue Specificity Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Cell Pathway Cell membrane; Multi-pass membrane protein.	Synonyms	
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled recefamily., Background olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	·	Observed Band	34kD
Function function:Odorant receptor .,similarity:Belongs to the G-protein coupled rece family., Olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory rece proteins are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated		Cell Pathway	Cell membrane; Multi-pass membrane protein.
family., Olfactory receptor family 4 subfamily D member 9(OR4D9) Homo sapiens Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory rece proteins are members of a large family of G-protein-coupled receptors (GPC arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated	Tissue Specificity	Tienes Conseilieite	
Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory receptors are members of a large family of G-protein-coupled receptors (GPC) arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormor receptors and are responsible for the recognition and G protein-mediated		rissue Specificity	function:Odorant receptor .,similarity:Belongs to the G-protein coupled receptor 1 family.,
the genome. The nomenclature assigned to the olfactory receptor genes an	Olfactory receptors interact with odorant molecules in the nose, to initiate a neuronal response that triggers the perception of a smell. The olfactory recept proteins are members of a large family of G-protein-coupled receptors (GPCR arising from single coding-exon genes. Olfactory receptors share a 7-transmembrane domain structure with many neurotransmitter and hormone		

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658







	proteins for this organism is independent of other organisms. [provided by RefSeq, Jul 2008],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images Western blot analysis of lysates from KB cells, primary 1 kDa antibody was diluted at 1:1000, 4°over night 180 --140 --100 --75 --60 --45 --35 --OR4D9 25 --15 --10 --

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658