



PACR Polyclonal Antibody

ng polypeptide type I receptor (PACAP type I
numan protein . at AA range: 80-160
endogenous levels of protein.
erol, and 0.02% sodium azide.
om rabbit antiserum by pe-specific immunogen.
00
ane protein.
pression in the lung, liver, thymus, spleen,
AP-27 and PACAP-38. The activity of this which activate adenylyl cyclase. May regulate luteinizing hormone, growth hormone, lamine. May play a role in spermatogenesis muscle relaxation and secretion in the ongs to the G-protein coupled receptor 2 dant in the brain, low expression in the lung,
dant in the brain, low expression in the lung, I placenta.,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



	biological actions of adenylate cyclase activating polypeptide 1 and is positively coupled to adenylate cyclase. Multiple alternatively spliced transcript variants encoding distinct isoforms have been identified. [provided by RefSeq, Dec 2010],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images