



DLG1 Polyclonal Antibody

Catalog No	BYab-07673
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB;ELISA
Gene Name	DLG1
Protein Name	Disks large homolog 1 (Synapse-associated protein 97) (SAP-97) (SAP97) (hDlg)
Immunogen	Synthesized peptide derived from part region of human protein AA range: 70-120
Specificity	DLG1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	99kD
Cell Pathway	Membrane ; Peripheral membrane protein . Basolateral cell membrane . Endoplasmic reticulum membrane . Cell junction, synapse, postsynaptic density . Cell junction, synapse . Cell membrane, sarcolemma . Apical cell membrane . Cell junction . Cytoplasm . Colocalizes with EPB41 at regions of intercellular contacts. Basolateral in epithelial cells (PubMed:12807908). May also associate with endoplasmic reticulum membranes. Mainly found in neurons soma, moderately found at postsynaptic densities (By similarity) .
Tissue Specificity	Abundantly expressed in atrial myocardium (at protein level). Expressed in lung fibroblasts, cervical epithelial and B-cells (at protein level). Expressed in the brain (at protein level) (PubMed:23676497). Widely expressed, with isoforms displaying different expression profiles.
Function	domain:The alternatively spliced domain I3 corresponding to amino acids (636-669) of isoform 4 is an EPB41 binding site mediating association to membranes in polarized and non-polarized cells.,domain:The L27 domain may regulate DLG1 self-association. The N-terminal alternatively spliced region is

Nanjing BYabscience technology Co.,Ltd



capable of binding several SH3 domains and also moderates the level of protein oligomerization. domain: The PDZ domains may also mediate association to membranes by binding to EPB41 and GPR124 together with the L27 domain that binds CASK and DLG2. function: Essential multidomain scaffolding protein required for normal development (By similarity). Recruits channels, receptors and signaling molecules to discrete plasma membrane domains in polarized cells. May play a role in adherens junction assembly, signal transduction, cell proliferation, synaptogenesis and lymphocyte activation. Regulates the ex

Background

This gene encodes a multi-domain scaffolding protein that is required for normal development. This protein may have a role in septate junction formation, signal transduction, cell proliferation, synaptogenesis and lymphocyte activation. Several alternatively spliced transcript variants encoding different isoforms have been described for this gene, but the full-length nature of some of the variants is not known. [provided by RefSeq, Feb 2011],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images