



DNMBP Polyclonal Antibody

| Catalog No | BYab-07675 |
|--------------------|---|
| Isotype | lgG |
| Reactivity | Human;Rat;Mouse; |
| Applications | WB;ELISA |
| Gene Name | DNMBP KIAA1010 |
| Protein Name | Dynamin-binding protein (Scaffold protein Tuba) |
| Immunogen | Synthesized peptide derived from part region of human protein AA range: 940-990 |
| Specificity | DNMBP Polyclonal Antibody detects endogenous levels of protein. |
| Formulation | Liquid in PBS containing 50% glycerol, and 0.02% sodium azide. |
| Source | Polyclonal, Rabbit,IgG |
| Purification | The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen. |
| Dilution | WB 1:500-2000 ELISA 1:5000-20000 |
| Concentration | 1 mg/ml |
| Purity | ≥90% |
| Storage Stability | -20°C/1 year |
| Synonyms | |
| Observed Band | 173kD |
| Cell Pathway | Cytoplasm . Golgi apparatus, Golgi stack . Cytoplasm, cytoskeleton . Cell junction, synapse . Cell junction . Localizes to the apical junction, colocalizes with TJP1 |
| Tissue Specificity | Detected in heart, brain, lung, liver, skeletal muscle, kidney and pancreas. |
| Function | function:Scaffold protein that links dynamin with actin-regulating proteins. May play a role in membrane trafficking between the cell surface and the Golgi.,similarity:Contains 1 BAR domain.,similarity:Contains 1 DH (DBL-homology) domain.,similarity:Contains 6 SH3 domains.,subcellular location:Localized to synapses and Golgi stacks.,subunit:Binds DNM1 via its N-terminal SH3 domains. The C-terminal SH3 domain binds a complex containing actin, tubulin, Hsp70 and actin-regulatory proteins, such as ENAH, EVL, WASL, WIRE, CR16, WAVE1 and NAP1L1.,tissue specificity:Detected in heart, brain, lung, liver, skeletal muscle, kidney and pancreas., |

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务,欢迎咨询



| Background | This gene encodes a protein belonging to the guanine nucleotide exchange factor family, and which regulates the configuration of cell junctions. It contains multiple binding sites for dynamin and thus links dynamin to actin regulatory proteins. Polymorphisms in this gene have been linked to Alzheimer's disease in some populations, though there are conflicting reports of such linkages in other populations. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Dec 2015], |
|---------------------------|--|
| matters needing attention | Avoid repeated freezing and thawing! |
| Usage suggestions | This product can be used in immunological reaction related experiments. For more information, please consult technical personnel. |

Products Images

Nanjing BYabscience technology Co.,Ltd