



S29A2 Polyclonal Antibody

Catalog No	BYab-07799
lsotype	lgG
Reactivity	Human;Mouse;Rat
Applications	WB;ELISA
Gene Name	SLC29A2 DER12 ENT2 HNP36
Protein Name	Equilibrative nucleoside transporter 2 (36 kDa nucleolar protein HNP36) (Delayed-early response protein 12) (Equilibrative nitrobenzylmercaptopurine riboside-insensitive nucleoside transporter) (Equil
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	S29A2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
• • •	
Concentration	1 mg/ml
Concentration	1 mg/ml ≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year 50kD Basolateral cell membrane ; Multi-pass membrane protein . Nucleus membrane ; Multi-pass membrane protein . Localized at the basolateral cell membrane in
Purity Storage Stability Synonyms Observed Band Cell Pathway	 ≥90% -20°C/1 year 50kD Basolateral cell membrane ; Multi-pass membrane protein . Nucleus membrane ; Multi-pass membrane protein . Localized at the basolateral cell membrane in polarized MDCK cells. Expressed in skeletal muscle, liver, lung, placenta, brain, heart, kidney and ovarian tissues.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	 ≥90% -20°C/1 year 50kD Basolateral cell membrane ; Multi-pass membrane protein . Nucleus membrane ; Multi-pass membrane protein . Localized at the basolateral cell membrane in polarized MDCK cells. Expressed in skeletal muscle, liver, lung, placenta, brain, heart, kidney and ovarian tissues. function:Mediates equilibrative transport of purine, pyrimidine nucleosides and the purine base hypoxanthine. Less sensitive than SLC29A1 to inhibition by nitrobenzylthioinosine (NBMPR), dipyridamole, dilazep and draflazine.,induction:By platelet derived growth factor (PDGF) and fibroblast growth factor (FGF).,similarity:Belongs to the SLC29A transporter family.,tissue

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务

24H 在线服务,欢迎咨询



physiologic processes through its effect on adenosine concentration at the cell surface (Griffiths et al., 1997 [PubMed 9396714]).[supplied by OMIM, Nov 2008],

matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images

Nanjing BYabscience technology Co.,Ltd