



HEY1 Polyclonal Antibody

Catalog No	BYab-07807
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	HEY1 BHLHB31 CHF2 HERP2 HESR1 HRT1
Protein Name	Hairy/enhancer-of-split related with YRPW motif protein 1 (Cardiovascular helix-loop-helix factor 2) (CHF-2) (Class B basic helix-loop-helix protein 31) (bHLHb31) (HES-related repressor protein 1) (Ha
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	HEY1 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	33kD
Cell Pathway	Nucleus .
Tissue Specificity	Expressed in the somitic mesoderm, the central nervous system, the kidney, the heart, nasal epithelium, and limbs.
Function	function:Downstream effector of Notch signaling which may be required for cardiovascular development. Transcriptional repressor which binds preferentially to the canonical E box sequence 5'-CACGTG-3'. Represses transcription by the cardiac transcriptional activators GATA4 and GATA6.,similarity:Belongs to the HEY family.,similarity:Contains 1 basic helix-loop-helix (bHLH) domain.,similarity:Contains 1 Orange domain.,subunit:Self-associates (By similarity). Interacts with HES1 and HEYL (By similarity). Interacts with HDAC1, NCOR1 and SIN3A (By similarity). Interacts with GATA4 and GATA6 (By similarity). Interacts with CCDC89/BOIP.,tissue specificity:Expressed in the somitic mesoderm, the central nervous system, the kidney, the heart, nasal

Nanjing BYabscience technology Co.,Ltd



epithelium, and limbs.,

Background

This gene encodes a nuclear protein belonging to the hairy and enhancer of split-related (HESR) family of basic helix-loop-helix (bHLH)-type transcriptional repressors. Expression of this gene is induced by the Notch and c-Jun signal transduction pathways. Two similar and redundant genes in mouse are required for embryonic cardiovascular development, and are also implicated in neurogenesis and somitogenesis. Alternative splicing results in multiple transcript variants. [provided by RefSeq, Jul 2008],


matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from A431 cells, primary antibody was diluted at 1:1000, 4° over night