



# HIC2 Polyclonal Antibody

<b>Catalog No</b>	BYab-07808
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	HIC2 HRG22 KIAA1020 ZBTB30
<b>Protein Name</b>	Hypermethylated in cancer 2 protein (Hic-2) (HIC1-related gene on chromosome 22 protein) (Hic-3) (Zinc finger and BTB domain-containing protein 30)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	HIC2 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	67kD
<b>Cell Pathway</b>	Nucleus.
<b>Tissue Specificity</b>	Highest levels in cerebellum.
<b>Function</b>	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,
<b>Background</b>	function:Transcriptional repressor.,similarity:Belongs to the krueppel C2H2-type zinc-finger protein family. Hic subfamily.,similarity:Contains 1 BTB (POZ) domain.,similarity:Contains 5 C2H2-type zinc fingers.,subunit:Interacts with CtBP.,tissue specificity:Highest levels in cerebellum.,
<b>matters needing attention</b>	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd



**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**