



BIRC2 Polyclonal Antibody

Catalog No	BYab-07853
Isotype	IgG
Reactivity	Human;Mouse
Applications	WB;ELISA
Gene Name	BIRC2 API1 IAP2 MIHB RNF48
Protein Name	Baculoviral IAP repeat-containing protein 2 (EC 6.3.2.-) (C-IAP1) (IAP homolog B) (Inhibitor of apoptosis protein 2) (IAP-2) (hIAP-2) (RING finger protein 48) (TNFR2-TRAF-signaling complex pro
Immunogen	Synthesized peptide derived from part region of human protein
Specificity	BIRC2 Polyclonal Antibody detects endogenous levels of protein.
Formulation	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
Dilution	WB 1:500-2000 ELISA 1:5000-20000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	
Observed Band	67kD
Cell Pathway	Cytoplasm. Nucleus. Agents that induce either the extrinsic or intrinsic apoptotic pathways promote its redistribution from the nuclear compartment to the cytoplasmic compartment. Associated with the midbody in telophase cells, and found diffusely in the nucleus of interphase cells.
Tissue Specificity	Present in many fetal and adult tissues. Mainly expressed in adult skeletal muscle, thymus, testis, ovary, and pancreas, low or absent in brain and peripheral blood leukocytes.
Function	function:Apoptotic suppressor. The BIR motifs region interacts with TNF receptor associated factors 1 and 2 (TRAF1 and TRAF2) to form an heteromeric complex, which is then recruited to the tumor necrosis factor receptor 2 (TNFR2).,similarity:Belongs to the IAP family.,similarity:Contains 1 CARD domain.,similarity:Contains 1 RING-type zinc finger.,similarity:Contains 3 BIR repeats.,subunit:Interacts with SMAC and with PRSS25; these interactions inhibit apoptotic suppressor activity.,tissue specificity:Present in many fetal and adult

Nanjing BYabscience technology Co.,Ltd



tissues. Mainly expressed in adult skeletal muscle, thymus, testis, ovary, and pancreas, low or absent in brain and peripheral blood leukocytes.,

Background

The protein encoded by this gene is a member of a family of proteins that inhibits apoptosis by binding to tumor necrosis factor receptor-associated factors TRAF1 and TRAF2, probably by interfering with activation of ICE-like proteases. This encoded protein inhibits apoptosis induced by serum deprivation and menadione, a potent inducer of free radicals. Alternatively spliced transcript variants encoding different isoforms have been found for this gene. [provided by RefSeq, Jan 2012],

matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images