



# DERL2 Polyclonal Antibody


<b>Catalog No</b>	BYab-07866
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human;Mouse
<b>Applications</b>	WB;ELISA
<b>Gene Name</b>	DERL2 DER2 FLANA CGI-101 SBBI53
<b>Protein Name</b>	Derlin-2 (Degradation in endoplasmic reticulum protein 2) (DERtrin-2) (Der1-like protein 2) (F-LAN-1) (F-LANa)
<b>Immunogen</b>	Synthesized peptide derived from part region of human protein
<b>Specificity</b>	DERL2 Polyclonal Antibody detects endogenous levels of protein.
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, and 0.02% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit antiserum by affinity-chromatography using epitope-specific immunogen.
<b>Dilution</b>	WB 1:500-2000 ELISA 1:5000-20000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	
<b>Observed Band</b>	26kD
<b>Cell Pathway</b>	Endoplasmic reticulum membrane ; Multi-pass membrane protein .
<b>Tissue Specificity</b>	Ubiquitous. Overexpressed in various hepatocarcinomas.
<b>Function</b>	function:Functional component of endoplasmic reticulum-associated degradation (ERAD) for misfolded luminal glycoproteins, but not that of misfolded nonglycoproteins. May act by forming a channel that allows the retrotranslocation of misfolded glycoproteins into the cytosol where they are ubiquitinated and degraded by the proteasome. May mediate the interaction between VCP and the degradation substrate. In contrast to DERL1, it is not involved in the degradation of MHC class I heavy chains following infection by cytomegaloviruses. May play a role in cell proliferation.,induction:Up-regulated in response to ER stress via the ERN1-XBP1 pathway of the unfolded protein response (UPR).,similarity:Belongs to the derlin family.,subunit:Forms homo- and heterooligomers with DERL3 and, to a lesser extent, with DERL1. Interacts with SELS, VCP and EDEM1. Mediates association between VCP and EDEM1, as

Nanjing BYabscience technology Co.,Ltd



<b>Background</b>	derlin 2(DERL2) Homo sapiens Proteins that are unfolded or misfolded in the endoplasmic reticulum (ER) must be refolded or degraded to maintain the homeostasis of the ER. DERL2 is involved in the degradation of misfolded glycoproteins in the ER (Oda et al., 2006 [PubMed 16449189]).[supplied by OMIM, Mar 2008],
<b>matters needing attention</b>	Avoid repeated freezing and thawing!
<b>Usage suggestions</b>	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## Products Images



Western blot analysis of lysates from K562 cells, primary antibody was diluted at 1:1000, 4° over night