




PLEK2 rabbit pAb

Catalog No BYab-07924 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name PLEK2 Protein Name PLEK2 Immunogen Synthesized peptide derived from human PLEK2 AA range: 5-55 Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation. similarity:Contains 1 DEP domain. similarity:Contains 2 PH domains. Background The protein encoded		
Reactivity Human; Mouse Applications WB Gene Name PLEK2 Protein Name PLEK2 Immunogen Synthesized peptide derived from human PLEK2 AA range: 5-55 Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix, Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation., similarity:Contains 1 DEP domains., similarity:Contains 2 PH domains. The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3 kinase. The encoded protein thein interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C10B), this gene shows an increase in expression in melanoma cellel spreading. In conjunction with complement component, a chain gervery as were as	Catalog No	BYab-07924
Applications Gene Name PLEK2 Protein Name PLEK2 Immunogen Synthesized peptide derived from human PLEK2 AA range: 5-55 Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix, Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation, similarity:Contains 1 DEP domain., similarity:Contains 2 PH domains. Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component, 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells a hading very as	Isotype	IgG
Gene Name PLEK2 Protein Name PLEK2 Immunogen Synthesized peptide derived from human PLEK2 AA range: 5-55 Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, 1, g subco	Reactivity	Human; Mouse
Protein Name PLEK2 Immunogen Synthesized peptide derived from human PLEK2 AA range: 5-55 Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, lgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix, Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation., similarity:Contains 1 DEP domain., similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, 1, generated and payeerye as the segment of the protein encoded protein in m	Applications	WB
Immunogen Synthesized peptide derived from human PLEK2 AA range: 5-55 Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Gene Name	PLEK2
Specificity This antibody detects endogenous levels of PLEK2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity 290% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domains.,similarity:Contains 2 PH domains. Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Protein Name	PLEK2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix, Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation., similarity:Contains 1 DEP domain., similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Immunogen	Synthesized peptide derived from human PLEK2 AA range: 5-55
Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix, Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation., similarity:Contains 1 DEP domain., similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Specificity	This antibody detects endogenous levels of PLEK2 at Human/Mouse
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.38% sodium azide.
using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains. Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Purification	, , , , , , , , , , , , , , , , , , , ,
Purity ≥90% Storage Stability -20°C/1 year Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Dilution	WB 1:500-2000
Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Concentration	1 mg/ml
Synonyms Pleckstrin-2 Observed Band 38kD Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Purity	≥90%
Observed Band Cell Pathway Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Storage Stability	-20°C/1 year
Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Synonyms	Pleckstrin-2
Cytoplasm, cytoskeleton. Tissue Specificity Cervix,Ovary, function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Observed Band	38kD
Function function:May help orchestrate cytoskeletal arrangement. Contribute to lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Cell Pathway	Cell projection, lamellipodium membrane; Peripheral membrane protein. Cytoplasm, cytoskeleton.
lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2 PH domains., Background The protein encoded by this gene associates with membrane-bound phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Tissue Specificity	Cervix,Ovary,
phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as	Function	lamellipodia formation.,similarity:Contains 1 DEP domain.,similarity:Contains 2
	Background	phosphatidylinositols generated by phosphatidylinositol 3-kinase. The encoded protein then interacts with the actin cytoskeleton to induce cell spreading. In conjunction with complement component 1, q subcomponent, B chain (C1QB), this gene shows an increase in expression in melanoma cells and may serve as

Nanjing BYabscience technology Co.,Ltd








matters needing attention

Avoid repeated freezing and thawing!

Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from MCF-7 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd