



GRAM2 rabbit pAb

Catalog No	BYab-07940
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	GRAMD2
Protein Name	GRAM2
Immunogen	Synthesized peptide derived from human GRAM2 AA range: 150-200
Specificity	This antibody detects endogenous levels of GRAM2 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.54% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	-20°C/1 year
Synonyms	GRAM domain-containing protein 2
Observed Band	38kD
Cell Pathway	Endoplasmic reticulum membrane ; Single-pass membrane protein . Cell membrane ; Peripheral membrane protein . Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Anchored at the ER, PM binding is mediated via GRAM domain and phosphatidylinositol lipid interaction. Localizes to distinct EPCS than GRAMD1A
Cell Pathway Tissue Specificity	Endoplasmic reticulum membrane ; Single-pass membrane protein . Cell membrane ; Peripheral membrane protein . Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Anchored at the ER, PM binding is mediated via GRAM domain and phosphatidylinositol lipid interaction.
	Endoplasmic reticulum membrane; Single-pass membrane protein. Cell membrane; Peripheral membrane protein. Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Anchored at the ER, PM binding is mediated via GRAM domain and phosphatidylinositol lipid interaction. Localizes to distinct EPCS than GRAMD1A
Tissue Specificity	Endoplasmic reticulum membrane; Single-pass membrane protein. Cell membrane; Peripheral membrane protein. Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Anchored at the ER, PM binding is mediated via GRAM domain and phosphatidylinositol lipid interaction. Localizes to distinct EPCS than GRAMD1A Brain, Hippocampus,
Tissue Specificity Function	Endoplasmic reticulum membrane; Single-pass membrane protein. Cell membrane; Peripheral membrane protein. Localizes to endoplasmic reticulum-plasma membrane contact sites (EPCS). Anchored at the ER, PM binding is mediated via GRAM domain and phosphatidylinositol lipid interaction. Localizes to distinct EPCS than GRAMD1A Brain, Hippocampus,

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from 3T3 cells, primary antibody was diluted at 1:1000, 4°over night

Nanjing BYabscience technology Co.,Ltd