



NXP20 rabbit pAb


Catalog No	BYab-07996
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	FAM114A1 NOXP20
Protein Name	NXP20
Immunogen	Synthesized peptide derived from human NXP20 AA range: 135-185
Specificity	This antibody detects endogenous levels of NXP20 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.110% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1:500-2000
Concentration	1 mg/ml
Purity	≥90%
Storage Stability	00004
Otorage Gtability	-20°C/1 year
Synonyms	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1)
	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein
Synonyms	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1)
Synonyms Observed Band	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1) 60kD
Synonyms Observed Band Cell Pathway	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1) 60kD Cytoplasm .
Synonyms Observed Band Cell Pathway Tissue Specificity	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1) 60kD Cytoplasm . Colon,Kidney,Salivary gland,Trachea, function:May play a role in neuronal cell development.,similarity:Belongs to the
Synonyms Observed Band Cell Pathway Tissue Specificity Function	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1) 60kD Cytoplasm . Colon,Kidney,Salivary gland,Trachea, function:May play a role in neuronal cell development.,similarity:Belongs to the
Synonyms Observed Band Cell Pathway Tissue Specificity Function Background matters needing	Protein NOXP20 (Nervous system overexpressed protein 20) (Protein FAM114A1) 60kD Cytoplasm . Colon,Kidney,Salivary gland,Trachea, function:May play a role in neuronal cell development.,similarity:Belongs to the FAM114 family.,

Nanjing BYabscience technology Co.,Ltd





Products Images



Western blot analysis of lysates from SW480 cells, primary antibody was diluted at 1:1000, 4°over night