



INSM2 rabbit pAb

Catalog No BYab-08019 Isotype IgG Reactivity Human; Mouse Applications WB Gene Name INSM2 IA6 Nbla106 Protein Name INSM2 Immunogen Synthesized peptide derived from human INSM2 AA range: 8-58 Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .		
Reactivity Human; Mouse Applications WB Gene Name INSM2 IA6 Nbla106 Protein Name INSM2 Immunogen Synthesized peptide derived from human INSM2 AA range: 8-58 Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Catalog No	BYab-08019
Applications WB Gene Name INSM2 IA6 Nbla106 Protein Name INSM2 Immunogen Synthesized peptide derived from human INSM2 AA range: 8-58 Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Isotype	IgG
Gene Name INSM2 IA6 Nbla106 Protein Name INSM2 Immunogen Synthesized peptide derived from human INSM2 AA range: 8-58 Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Reactivity	Human; Mouse
Protein Name INSM2 Immunogen Synthesized peptide derived from human INSM2 AA range: 8-58 Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Applications	WB
Immunogen Synthesized peptide derived from human INSM2 AA range: 8-58 Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Gene Name	INSM2 IA6 Nbla106
Specificity This antibody detects endogenous levels of INSM2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Protein Name	INSM2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Immunogen	Synthesized peptide derived from human INSM2 AA range: 8-58
SourcePolyclonal, Rabbit,IgGPurificationThe antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.DilutionWB 1:500-2000Concentration1 mg/mlPurity≥90%Storage Stability-20°C/1 yearSynonymsInsulinoma-associated protein 2 (Zinc finger protein IA-6)Observed Band60kDCell PathwayCytoplasm . Nucleus .	Specificity	This antibody detects endogenous levels of INSM2 at Human/Mouse
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.133% sodium azide.
Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Purification	, , , , , , , , , , , , , , , , , , , ,
Purity ≥90% Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Dilution	WB 1:500-2000
Storage Stability -20°C/1 year Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Concentration	1 mg/ml
Synonyms Insulinoma-associated protein 2 (Zinc finger protein IA-6) Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Purity	≥90%
Observed Band 60kD Cell Pathway Cytoplasm . Nucleus .	Storage Stability	-20°C/1 year
Cell Pathway Cytoplasm . Nucleus .	Synonyms	Insulinoma-associated protein 2 (Zinc finger protein IA-6)
	Observed Band	60kD
	Cell Pathway	Cytoplasm . Nucleus .
extent, in brain, lung and spleen. In the pancreas, expressed in islet cells, including insulin- and glucagon-producing alpha- and beta-cells, but not in acina	Tissue Specificity	including insulin- and glucagon-producing alpha- and beta-cells, but not in acinar cells (at protein level). Detected in adrenal glands, particularly in the deeper layer
Function function: May function as a growth suppressor or tumor suppressor in liver cells and in certain neurons., similarity: Contains 5 C2H2-type zinc fingers.,	Function	function:May function as a growth suppressor or tumor suppressor in liver cells and in certain neurons.,similarity:Contains 5 C2H2-type zinc fingers
Background	Background	, , , ,
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd