



MMP21 rabbit pAb

Function cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,		
Reactivity Human; Mouse Applications WB Gene Name MMP21 Protein Name MMP21 Immunogen Synthesized peptide derived from human MMP21 AA range: 425-475 Specificity This antibody detects endogenous levels of MMP21 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc lon per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thin inhibiting the enzyme. The dissociation of the cysteine from	Catalog No	BYab-08055
Applications WB Gene Name MMP21 Protein Name MMP21 Immunogen Synthesized peptide derived from human MMP21 AA range: 425-475 Specificity This antibody detects endogenous levels of MMP21 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine-switch motif binds the catalytic zinc ion, unsuring hipheral-taintypsin, PTW. The precursor is cleaved by a furine endopeptidase, similarity:Belongs to the peptidase M10A family, similarity Contains 4 hemospoxin-like domains. Issue specificity; Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.	Isotype	IgG
Gene Name MMP21 Protein Name MMP21 Immunogen Synthesized peptide derived from human MMP21 AA range: 425-475 Specificity This antibody detects endogenous levels of MMP21 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme, function: May have an important and specific function in tumor progression	Reactivity	Human; Mouse
Protein Name MMP21 Immunogen Synthesized peptide derived from human MMP21 AA range: 425-475 Specificity This antibody detects endogenous levels of MMP21 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme, function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin, plancenta, brain and pe	Applications	WB
Immunogen Synthesized peptide derived from human MMP21 AA range: 425-475 Specificity This antibody detects endogenous levels of MMP21 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. Introtion:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitypsin, PTM:The precursor is cleaved by a furin endopeptidase, similarity:Belongs to the peptidase M10A family, similarity:Contains 4 hemopexin-like domains, lissue specificity:Identified in fetal brain, kidney	Gene Name	MMP21
Specificity This antibody detects endogenous levels of MMP21 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zine ion per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme. function. May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitypsin, PTM:The precursor is cleaved by a furin endopeptidase, similarity:Belongs to the peptidase M10A family, similarity:Contains 4 hemopexin-like domains, Lissue specificity:Identified in fetal brain, kidney and liver. In adu	Protein Name	MMP21
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide. Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted. Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family, similarity.Contains 4 hemopex.in-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.	Immunogen	Synthesized peptide derived from human MMP21 AA range: 425-475
Source Polyclonal, Rabbit, IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit. cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme, function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase, similarity:Belongs to the peptidase M10A family, similarity:Contains 4 hemopexin-like domains "tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.	Specificity	This antibody detects endogenous levels of MMP21 at Human/Mouse
Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium, domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme, function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin, PTM:The precursor is cleaved by a furin endopeptidase, similarity:Contains 4 hemopexin-like domains, tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.169% sodium azide.
Dilution WB 1:500-2000 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme, function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidasesimilarity:Centains 4 hemopexin-like domains, tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase, similarity:Belongs to the peptidase M10A family., similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Purification	
Purity ≥90% Storage Stability -20°C/1 year Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Dilution	WB 1:500-2000
Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted. Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit, cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme. function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family, similarity:Contains 4 hemopexin-like domains., tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Concentration	1 mg/ml
Synonyms Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24) Observed Band 60kD Cell Pathway Secreted. Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Purity	≥90%
Cell Pathway Secreted . Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Storage Stability	-20°C/1 year
Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Function cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Synonyms	Matrix metalloproteinase-21 (MMP-21) (EC 3.4.24)
Tissue Specificity Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Observed Band	60kD
kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines. Cofactor:Binds 1 zinc ion per subunit.,cofactor:Calcium.,domain:The conserved cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Cell Pathway	Secreted .
cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in various cancer cell lines.,	Tissue Specificity	kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as
AL VY MAN I I	Function	cysteine present in the cysteine-switch motif binds the catalytic zinc ion, thus inhibiting the enzyme. The dissociation of the cysteine from the zinc ion upon the activation-peptide release activates the enzyme.,function:May have an important and specific function in tumor progression and embryogenesis. Cleaves alpha-1-antitrypsin.,PTM:The precursor is cleaved by a furin endopeptidase.,similarity:Belongs to the peptidase M10A family.,similarity:Contains 4 hemopexin-like domains.,tissue specificity:Identified in fetal brain, kidney and liver. In adult tissues found primarily in ovary, kidney, liver, lung, placenta, brain and peripheral blood leukocytes. Expressed as well in

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Background	This gene encodes a member of the matrix metalloproteinase family. Proteins in this family are involved in the breakdown of extracellular matrix for both normal physiological processes, such as embryonic development, reproduction, and tissue remodeling, and disease processes, such as asthma and tumor metastasis. The encoded protein may play an important role in embryogenesis, particularly in neuronal cells, as well as in lymphocyte development and survival. [provided by RefSeq, May 2013],
matters needing attention	Avoid repeated freezing and thawing!
Usage suggestions	This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658