



# ZNT5 rabbit pAb

<b>Catalog No</b>	BYab-08196
<b>Isotype</b>	IgG
<b>Reactivity</b>	Human; Mouse
<b>Applications</b>	WB
<b>Gene Name</b>	SLC30A5 ZNT5 ZNTL1 ZTL1 UNQ863/PRO1879
<b>Protein Name</b>	ZNT5
<b>Immunogen</b>	Synthesized peptide derived from human ZNT5 AA range: 106-156
<b>Specificity</b>	This antibody detects endogenous levels of ZNT5 at Human/Mouse
<b>Formulation</b>	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.311% sodium azide.
<b>Source</b>	Polyclonal, Rabbit,IgG
<b>Purification</b>	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
<b>Dilution</b>	WB 1:500-2000
<b>Concentration</b>	1 mg/ml
<b>Purity</b>	≥90%
<b>Storage Stability</b>	-20°C/1 year
<b>Synonyms</b>	Zinc transporter 5 (ZnT-5) (Solute carrier family 30 member 5) (ZnT-like transporter 1) (hZTL1)
<b>Observed Band</b>	85kD
<b>Cell Pathway</b>	Golgi apparatus, trans-Golgi network membrane ; Multi-pass membrane protein . Perimeter of granules, localizes to the brush border membrane of the enterocyte. Concentrated in early compartments of the secretory pathway such as COPII-coated vesicles.
<b>Tissue Specificity</b>	Ubiquitously expressed. Highly expressed in pancreas, liver and kidney. Expressed abundantly in insulin-containing beta cells, undetectable in other endocrine cell types including glucagon-secreting alpha cells and most acinar cells.
<b>Function</b>	function:Functions as a zinc transporter. May be a transporter of zinc into beta cells in order to form insulin crystals. Partly regulates cellular zinc homeostasis. Required with ZNT7 for the activation of zinc-requiring enzymes, alkaline phosphatases (ALPs). Transports zinc into the lumens of the Golgi apparatus and vesicular compartments where ALPs locate, thus, converting apoALPs to holoALPs. Required with ZNT6 and ZNT7 for the activation of TNAP.,induction:Increased intracellular zinc level, resulting from extracellular zinc

Nanjing BYabscience technology Co.,Ltd



supplementation, do not induce any up- or down-regulation of gene expression. Up-regulated by zinc depletion.,miscellaneous:The histidine-rich loop is essential for ZNT5 and ZNT6 heterooligomeric complexes to activate TNAP.,similarity:Belongs to the cation diffusion facilitator (CDF) transporter (TC 2.A.4) family. SLC30A subfamily.,subcellular location:Perimeter

**Background**

This gene encodes a member of the SLC30A/ZnT family of zinc transporter proteins. ZnT proteins mediate both cellular zinc efflux and zinc sequestration into membrane-bound organelles. The encoded protein plays a role in the early secretory pathway as a heterodimer with zinc transporter 6, and may also regulate zinc sequestration into secretory granules of pancreatic beta cells. Alternatively spliced transcript variants encoding multiple isoforms have been observed for this gene, and a pseudogene of this gene is located on the long arm of chromosome 19. [provided by RefSeq, Oct 2011],


**matters needing attention**

Avoid repeated freezing and thawing!

**Usage suggestions**

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

**Products Images**



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night