



ZBED5 rabbit pAb

Catalog No	BYab-08290
Isotype	IgG
Reactivity	Human;Rat;Mouse;
Applications	WB
Gene Name	ZBED5
Protein Name	ZBED5
Immunogen	Synthesized peptide derived from human ZBED5 AA range: 521-571
Specificity	This antibody detects endogenous levels of ZBED5 at Human
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration Purity	1 mg/ml ≥90%
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year nucleoplasm,cytoplasm, miscellaneous:May be derived from an ancient transposon that has lost its ability
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year nucleoplasm,cytoplasm, miscellaneous:May be derived from an ancient transposon that has lost its ability to translocate.,similarity:Contains 1 BED-type zinc finger., This gene is unusual in that its coding sequence is mostly derived from Charlie-like DNA transposon; however, it does not appear to be an active DNA transposon as it is not flanked by terminal inverted repeats. The encoded protein is conserved among the mammalian Laurasiatheria branch. Multiple alternatively spliced variants, encoding the same protein, have been identified. [provided by

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

Products Images



Western blot analysis of lysates from HpeG2 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd