



RCAN3 rabbit pAb

Catalog No	BYab-08431
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	RCAN3 DSCR1L2
Protein Name	RCAN3
Immunogen	Synthesized peptide derived from human RCAN3 AA range: 86-136
Specificity	This antibody detects endogenous levels of RCAN3 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concentration	9
Purity	≥90%
	-
Purity	≥90%
Purity Storage Stability	≥90%
Purity Storage Stability Synonyms	≥90%
Purity Storage Stability Synonyms Observed Band	≥90% -20°C/1 year
Purity Storage Stability Synonyms Observed Band Cell Pathway	≥90% -20°C/1 year intracellular,cytoplasm, Highest expression in heart, skeletal muscle kidney, liver and peripheral blood leukocytes. Lower expression in all other tissues. function:Inhibits calcineurin-dependent transcriptional responses by binding to the catalytic domain of calcineurin A. Could play a role during central nervous system developmentsimilarity:Belongs to the RCAN familytissue specificity:Highest
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	≥90% -20°C/1 year intracellular,cytoplasm, Highest expression in heart, skeletal muscle kidney, liver and peripheral blood leukocytes. Lower expression in all other tissues. function:Inhibits calcineurin-dependent transcriptional responses by binding to the catalytic domain of calcineurin A. Could play a role during central nervous system development.,similarity:Belongs to the RCAN family.,tissue specificity:Highest expression in heart, skeletal muscle kidney, liver and peripheral blood leukocytes.
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	≥90% -20°C/1 year intracellular,cytoplasm, Highest expression in heart, skeletal muscle kidney, liver and peripheral blood leukocytes. Lower expression in all other tissues. function:Inhibits calcineurin-dependent transcriptional responses by binding to the catalytic domain of calcineurin A. Could play a role during central nervous system development.,similarity:Belongs to the RCAN family.,tissue specificity:Highest expression in heart, skeletal muscle kidney, liver and peripheral blood leukocytes.

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658





国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.





Western blot analysis of lysates from A549 cells, primary antibody was diluted at 1:1000, 4° over night

Nanjing BYabscience technology Co.,Ltd