



ACOT6 rabbit pAb

Catalog No	BYab-08551
Isotype	IgG
Reactivity	Human; Mouse
Applications	WB
Gene Name	ACOT6 C14orf42
Protein Name	ACOT6
Immunogen	Synthesized peptide derived from human ACOT6 AA range: 116-166
Specificity	This antibody detects endogenous levels of ACOT6 at Human/Mouse
Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
Source	Polyclonal, Rabbit,IgG
Purification	The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.
Dilution	WB 1: 500-2000
Concentration	1 mg/ml
Concontitution	-
Purity	≥90%
	≥90% -20°C/1 year
Purity	
Purity Storage Stability	
Purity Storage Stability Synonyms	
Purity Storage Stability Synonyms Observed Band	-20°C/1 year [Isoform 1]: Peroxisome . Localization to the peroxisome is uncertain since the potential C-terminal peroxisome targeting signal found in the mouse ortholog is not perfectly conserved; [Isoform 2]: Cytoplasm . Recombinant N-terminally
Purity Storage Stability Synonyms Observed Band Cell Pathway	-20°C/1 year [Isoform 1]: Peroxisome . Localization to the peroxisome is uncertain since the potential C-terminal peroxisome targeting signal found in the mouse ortholog is not perfectly conserved; [Isoform 2]: Cytoplasm . Recombinant N-terminally GFP-tagged protein localizes to the cytosol caution:May lack catalytic activity, even though the active site residues are
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity	-20°C/1 year [Isoform 1]: Peroxisome . Localization to the peroxisome is uncertain since the potential C-terminal peroxisome targeting signal found in the mouse ortholog is not perfectly conserved; [Isoform 2]: Cytoplasm . Recombinant N-terminally GFP-tagged protein localizes to the cytosol caution:May lack catalytic activity, even though the active site residues are conserved.,function:Unknown.,similarity:Belongs to the C/M/P thioester hydrolase
Purity Storage Stability Synonyms Observed Band Cell Pathway Tissue Specificity Function	-20°C/1 year [Isoform 1]: Peroxisome . Localization to the peroxisome is uncertain since the potential C-terminal peroxisome targeting signal found in the mouse ortholog is not perfectly conserved; [Isoform 2]: Cytoplasm . Recombinant N-terminally GFP-tagged protein localizes to the cytosol caution:May lack catalytic activity, even though the active site residues are conserved.,function:Unknown.,similarity:Belongs to the C/M/P thioester hydrolase

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658



国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务,欢迎咨询



Usage suggestions

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

