



## MORN2 rabbit pAb

Catalog No       BYab-08589         Isotype       IgG         Reactivity       Human; Mouse         Applications       IHC;IF         Gene Name       MORN2 MOPT         Protein Name       MORN2         Immunogen       Synthesized peptide derived from human MORN2 AA range: 25-75         Specificity       This antibody detects endogenous levels of MORN2 at Human/Mouse         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       IHC-p 1: 50-200. IF 1:50-200         Concentration       1 mg/ml         Purity       290%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermogenesis. Appears to migrate to the nucleus during the final step of spermogenesis. Appears to migrate to the nucleus during the final step of spermogenesis.         Background       Avoid repeated freezing and thawing!		
Reactivity Human; Mouse  Applications IHC;IF  Gene Name MORN2 MOPT  Protein Name MORN2  Immunogen Synthesized peptide derived from human MORN2 AA range: 25-75  Specificity This antibody detects endogenous levels of MORN2 at Human/Mouse  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution IHC-p 1: 50-200. IF 1:50-200  Concentration 1 mg/ml  Purity 290%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis . Appears to migrate to the nucleus during the final step of spermiogenesis . Appears to migrate to the nucleus during the final step of spermiogenesis .  Background  matters needing Avoid repeated freezing and thawing!	Catalog No	BYab-08589
Applications IHC;IF Gene Name MORN2 MOPT Protein Name MORN2 Immunogen Synthesized peptide derived from human MORN2 AA range: 25-75 Specificity This antibody detects endogenous levels of MORN2 at Human/Mouse Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide. Source Polyclonal, Rabbit,IgG Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen. Dilution IHC-p 1: 50-200. IF 1:50-200 Concentration 1 mg/ml Purity ≥90% Storage Stability -20°C/1 year Synonyms Observed Band Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis . Appears to migrate to the nucleus during the final step of spermiogenesis . Appears to migrate to the nucleus during the final step of spermiogenesis .  Background matters needing Avoid repeated freezing and thawing!	Isotype	IgG
Gene Name MORN2 MOPT  Protein Name MORN2  Immunogen Synthesized peptide derived from human MORN2 AA range: 25-75  Specificity This antibody detects endogenous levels of MORN2 at Human/Mouse  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution IHC-p 1: 50-200. IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis.  Background  matters needing Avoid repeated freezing and thawing!	Reactivity	Human; Mouse
Protein Name         MORN2           Immunogen         Synthesized peptide derived from human MORN2 AA range: 25-75           Specificity         This antibody detects endogenous levels of MORN2 at Human/Mouse           Formulation         Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.           Source         Polyclonal, Rabbit, IgG           Purification         The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.           Dilution         IHC-p 1: 50-200. IF 1:50-200           Concentration         1 mg/ml           Purity         ≥90%           Storage Stability         -20°C/1 year           Synonyms         Observed Band           Cell Pathway         Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis . Appears to migrate to the nucleus during the final step of spermiogenesis . Appears to migrate to the nucleus during the final step of spermiogenesis .           Tissue Specificity         Function         similarity:Contains 2 MORN repeats.,           Background         Avoid repeated freezing and thawing!	Applications	IHC;IF
Immunogen       Synthesized peptide derived from human MORN2 AA range: 25-75         Specificity       This antibody detects endogenous levels of MORN2 at Human/Mouse         Formulation       Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.         Source       Polyclonal, Rabbit, IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       IHC-p 1: 50-200. IF 1:50-200         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis Appears to migrate to the nucleus during the final step of spermiogenesis.         Function       similarity:Contains 2 MORN repeats.,         Background       Avoid repeated freezing and thawing!	Gene Name	MORN2 MOPT
Specificity This antibody detects endogenous levels of MORN2 at Human/Mouse  Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit,IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution IHC-p 1: 50-200. IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis.  Background  matters needing Avoid repeated freezing and thawing!	Protein Name	MORN2
Formulation Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.  Source Polyclonal, Rabbit, IgG  Purification The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution IHC-p 1: 50-200. IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis.  Tissue Specificity  Function similarity:Contains 2 MORN repeats.,  Background  Matters needing Avoid repeated freezing and thawing!	Immunogen	Synthesized peptide derived from human MORN2 AA range: 25-75
Source       Polyclonal, Rabbit,IgG         Purification       The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.         Dilution       IHC-p 1: 50-200. IF 1:50-200         Concentration       1 mg/ml         Purity       ≥90%         Storage Stability       -20°C/1 year         Synonyms       Observed Band         Cell Pathway       Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis.         Tissue Specificity         Function       similarity:Contains 2 MORN repeats.,         Background       Avoid repeated freezing and thawing!	Specificity	This antibody detects endogenous levels of MORN2 at Human/Mouse
Purification  The antibody was affinity-purified from rabbit serum by affinity-chromatography using specific immunogen.  Dilution  IHC-p 1: 50-200. IF 1:50-200  Concentration  1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway  Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermlogenesis  Tissue Specificity  Function  similarity:Contains 2 MORN repeats.,  Background  Matters needing  Avoid repeated freezing and thawing!	Formulation	Liquid in PBS containing 50% glycerol, 0.5% BSA and 0.02% sodium azide.
using specific immunogen.  Dilution IHC-p 1: 50-200. IF 1:50-200  Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis  Tissue Specificity  Function similarity:Contains 2 MORN repeats.,  Background  Matters needing Avoid repeated freezing and thawing!	Source	Polyclonal, Rabbit,IgG
Concentration 1 mg/ml  Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis.  Tissue Specificity  Function similarity:Contains 2 MORN repeats.,  Background  Matters needing Avoid repeated freezing and thawing!	Purification	
Purity ≥90%  Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis  Tissue Specificity  Function similarity:Contains 2 MORN repeats.,  Background  matters needing Avoid repeated freezing and thawing!	Dilution	IHC-p 1: 50-200. IF 1:50-200
Storage Stability -20°C/1 year  Synonyms  Observed Band  Cell Pathway Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis  Tissue Specificity  Function similarity:Contains 2 MORN repeats.,  Background  Avoid repeated freezing and thawing!	Concentration	1 mg/ml
Synonyms  Observed Band  Cell Pathway  Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis  Tissue Specificity  Function  similarity:Contains 2 MORN repeats.,  Background  Matters needing  Avoid repeated freezing and thawing!	Purity	≥90%
Observed Band  Cell Pathway  Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis.  Tissue Specificity  Function  similarity:Contains 2 MORN repeats.,  Background  Avoid repeated freezing and thawing!	Storage Stability	-20°C/1 year
Cell Pathway  Cytoplasmic vesicle, secretory vesicle, acrosome . Nucleus . Associated with the acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis  Tissue Specificity  Function  similarity:Contains 2 MORN repeats.,  Background  Matters needing  Avoid repeated freezing and thawing!	Synonyms	
acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to the nucleus during the final step of spermiogenesis  Tissue Specificity  Function similarity:Contains 2 MORN repeats.,  Background  matters needing Avoid repeated freezing and thawing!	Observed Band	
Function similarity:Contains 2 MORN repeats.,  Background  matters needing Avoid repeated freezing and thawing!	Cell Pathway	acrosome during post-meiotic stages of spermatogenesis. Appears to migrate to
Background  matters needing Avoid repeated freezing and thawing!	Tissue Specificity	
matters needing Avoid repeated freezing and thawing!	Function	similarity:Contains 2 MORN repeats.,
	Background	
	matters needing attention	Avoid repeated freezing and thawing!

Nanjing BYabscience technology Co.,Ltd

网址: www.njbybio.com 官方热线: 025-5229-8998 监督电话: 15950492658




国内优质抗体供应商 精准的 WB 检测服务 24H 在线服务, 欢迎咨询



**Usage suggestions** 

This product can be used in immunological reaction related experiments. For more information, please consult technical personnel.

## **Products Images**



Immunohistochemical analysis of paraffin-embedded human Gastric adenocarcinoma. 1, Antibody was diluted at 1:200(4° overnight). 2, Tris-EDTA,pH9.0 was used for antigen retrieval. 3,Secondary antibody was diluted at 1:200(room temperature, 45min).

Nanjing BYabscience technology Co.,Ltd